

Drought Stress Amelioration in Sesame Cultivars: Unraveling the Combined Potential of MnSO_4 Seed Treatment and Foliar Nutrition

Ghasem Najafi , Mohammad Javad Zarea* , Arash Fazeli , Batool Zarei 

Department of Agronomy and Crop Breeding, Faculty of Agriculture, Ilam University, Ilam, Iran

ARTICLE INFO

Original paper

Article history:

Received: x Month 202x

Revised: x Month 202x

Accepted: x Month 202x

Keywords:

Antioxidant enzymes

Manganese sulfate

Priming

Sesame

Water deficiency

ABSTRACT

Water scarcity is a critical factor that restricts crop production, and projections indicate that its severity will escalate in the future due to the impacts of global warming. The objective of this research was to assess the role of MnSO_4 in improving water deficit tolerance in two sesame cultivars, Dashtestan2 and Naz Takshakheh. A factorial pot study was conducted based on randomized complete block design with three replications at the Research Greenhouse of the Faculty of Agriculture, University of Ilam in 2019. $\text{MnSO}_4 \cdot 7\text{H}_2\text{O}$ was utilized via seed priming, foliar application, or a combination of both methods. Three different water level regimes were established: 75% field capacity (FC) representing well-watered conditions, 50% FC indicating mild water stress, and 25% FC denoting severe water stress. The results of experiment indicated that cultivar Dashtestan2 produced a greater seed yield compared to the Naz Takshakheh cultivar, under both well-watered and water-deficient conditions. Foliar application was more effective than seed priming in promoting proline accumulation and enhancing the activities of catalase and ascorbate peroxidase enzymes. Plants treated with both seed priming and foliar application with $\text{MnSO}_4 \cdot 7\text{H}_2\text{O}$ demonstrated enhanced antioxidant enzyme activity and higher seed yield. Based on the results, the application of manganese sulfate through seed priming, foliar application, and a combination of both techniques led to a notable increase in seed yield for both sesame cultivars facing water deficit stress. The Dashtestan2 cultivar showed especially significant improvements in seed yield, which can be attributed to increased activity of antioxidant enzymes and osmotic regulators. Among the various methods for applying manganese sulfate, the combination of seed priming and foliar spraying demonstrated superior effectiveness in enhancing tolerance to water deficit stress.

DOI: [10.22126/ATIC.2026.12043.1212](https://doi.org/10.22126/ATIC.2026.12043.1212)

© The Author(s) 202x. Published by Razi University



1. Introduction

Sesame (*Sesamum indicum* L.) is a vital oilseed valued for its various industrial, pharmaceutical, and nutritional uses (Kermani *et al.*, 2019; Hussain *et al.*, 2023). The seeds are rich in protein, oil, and natural antioxidants such as sesamin, sesamol, and tocopherol (Balouchi *et al.*, 2023). Sesame is typically cultivated in arid and semi-arid regions worldwide. Plants are exposed to various environmental stresses during growth and development under natural and agricultural conditions. Among these, drought is one of the most severe environmental stresses that severely affects crop growth and productivity, especially in arid and semi-arid regions such as Iran, where annual average rainfall is about 250 mm (Hamarash *et al.*,

2022). Drought stress disrupts growth by causing various morphological, physiological, biochemical, and molecular changes, which ultimately affect the crop's final yield (Rasheed and Malik, 2022). The response of plants to drought stress depends on growth stage, species, the severity and duration of the drought (Gray and Brady, 2016). Water deficiency reduces photosynthesis by reducing leaf area index, stomata conductance, and photosynthetic pigments concentration. Under water-deficient conditions, plant growth and yield are reduced due to the limited production of photosynthetic materials (Dawood *et al.*, 2019). Water deficiency, the most limiting stress in agriculture, can severely impact both seed quality and yield. Therefore, investigating how plants cope with

* Corresponding author.

E-mail address: zareamj@yahoo.com

water limitation is highly valuable and should be prioritized in the near future, particularly in arid and semi-arid environments such as Iran (Sobhanian et al., 2020). Plants respond to water deficiency by increasing the activity of antioxidant defense systems to eliminate reactive oxygen free radicals (Kar, 2011) and accumulating organic osmolytes to maintain osmotic balance (Blum, 2017). When plants experience drought stress, they activate a range of antioxidant defenses to mitigate the harmful effects of oxidative stress. These include enzymes like catalase (CAT), superoxide dismutase (SOD), peroxidase (POX), ascorbate peroxidase (APX), and non-enzymatic antioxidants like glutathione and ascorbate (Chen et al., 2016).

Compatible compounds such as proline, glycine betaine or polyamines, as well as sugars, are produced as osmotic protectors in plants during stress conditions (Hasanuzzaman et al., 2019). Most plants respond to stress conditions such as drought, salinity, nutrient deficiency, and high temperatures by increasing proline levels (Jogawat, 2019). Since proline is an important osmotic regulator, the accumulation of proline increases resistance to water deficiency in plants (Su et al., 2018). Glycine betaine, along with other osmotic regulators, is commonly accumulated in plants during stress conditions (Hernandez-Leon and Valenzuela-Soto, 2023). Many strategies exist to enhance the growth and development of plant species under both stress and non-stress conditions. One effective approach is seed priming, a chemical treatment applied before sowing. This method promotes early seedling emergence and accelerates the emergence rate, even in extreme weather conditions (Ghosh et al., 2023). During priming, seeds are soaked in water or in various solutions, including organic and inorganic salts, within a controlled environment. After soaking, the seeds are dried before sowing. This process helps regulate germination by managing the moisture and temperature of the seeds (Diya et al., 2024). Priming allows seeds to absorb sufficient water for germination using less moisture than is usually required for natural germination (Farooq et al., 2019). Many organic and mineral substances, hormones, and plant extracts have proven effective as priming agents in various crops against abiotic stresses, including drought and salinity stresses (Amritha et al., 2021). Manganese sulfate is a key basic manganese salt, accounting for nearly 80% of the world's manganese products, which are produced

using either manganese sulfate or manganese sulfate solutions. Additionally, manganese sulfate has numerous applications in both industry and agriculture (Li et al., 2023). Priming maize seeds with a MnSO_4 solution enhanced growth, grain manganese content, and seed yield. Moreover, increasing the concentration of the priming solution to 0.2% MnSO_4 for 12 hours resulted in a linear increase in both seed yield and grain manganese content (Muhammad et al., 2015).

This study aims to investigate the efficacy of manganese sulfate ($\text{MnSO}_4 \cdot 7\text{H}_2\text{O}$) in enhancing drought tolerance in two sesame cultivars (Dashtestan2 and Naz Takshakheh) under varying water deficit conditions. Specifically, the research evaluates the impact of different application methods (seed priming, foliar spraying, and their combination) on antioxidant enzyme activity (CAT and APX), proline accumulation, and seed yield at three irrigation levels (75%, 50%, and 25% of field capacity). Additionally, the study compares cultivar-specific responses to water stress and identifies the most effective MnSO_4 application strategy for mitigating drought-induced yield losses. The findings will provide critical insights into optimizing sesame production under water-scarce conditions, contributing to sustainable agriculture in arid and semi-arid regions.

2. Materials and methods

A factorial experiment was carried out based on randomized complete block design with three replications to assess the impact of seed priming and foliar application of $\text{MnSO}_4 \cdot 7\text{H}_2\text{O}$ on the growth and yield of sesame under water deficiency conditions. The research was conducted at the greenhouse of the Faculty of Agriculture, the University of Ilam, in December 2019 (min and max temperatures averaged 24°C and 34°C, respectively). Three experimental factors were used including two sesame cultivars (Dashtestan2 and Naz Takshakheh), different application types including seed priming (unprimed seeds and seed primed) and foliar application (control and treated), and the third factor including three level of water stress (well-watered (75% FC), mild water deficiency (50% FC) and severe water deficiency (25% FC). Seeds from two sesame varieties, Dashtestan2 and Naz Takshakheh, were acquired from the Seed and Plant Improvement Institute located in Karaj, Iran (Table 1).

Table 1. Properties of the studied cultivars

Cultivars	Branching	Seed color	1000 seed weight (g)	Response to water deficiency	Plant height (cm)
Dashtestan2	Several branches	Light brown	4	Tolerant	142-145
Naz TakShaskheh	Single branches	Cream	2.6-2.9	Sensitive	100-120

The experimental units consisted of 72 plastic pots, each with dimensions of 25 cm in diameter and height, filled with 7 kg of field soil. To ensure proper drainage, four holes were drilled at the bottom of each pot, and equal-weight pebbles were added to all of them. Before planting, the soil's texture, pH, and electrical conductivity (EC) were measured at the soil laboratory at Ilam University. Following the soil analysis, fertilizers including phosphorus (triple superphosphate), nitrogen (urea), and potash fertilizers were incorporated into the soil in the pots (Table 2).

Table 2. Soil physicochemical properties

Texture	E.C. (dS m ⁻¹)	pH	O.C. (%)	N (%)	Mn (mg kg ⁻¹)	P (mg kg ⁻¹)	K (mg kg ⁻¹)
Clay loam	0.3	7	1.2	0.12	0.08	8.5	420

The process of disinfecting seeds prior to planting consisted of multiple stages. Initially, the seeds were treated with 70% ethanol for a duration of 30 seconds, then immersed in a 5% sodium hypochlorite solution for five minutes. The seeds were then thoroughly washed multiple times with distilled water to eliminate any remaining sodium hypochlorite. Ten seeds were planted in each pot at a depth of 1 cm. After germination, five plants were retained in each pot. Sesame plant growth was carried out in a greenhouse with average day and night temperatures of 34°C and 24°C, respectively, relative humidity of 45-60%, and a light period of 16 hours. Pots were regularly irrigated prior to the onset of the water deficiency treatment to maintain soil moisture at 75% of FC, which is 40% for this soil type. Irrigation treatments began one month after plant emergence. The weights of each pot for the three treatments—75%, 50%, and 25% FC were determined to be 1.10 kg, 9.4 kg, and 8.7 kg, respectively. A pot weighing system was used to determine irrigation time, and irrigation was performed when the pot weight for each treatment reached a predetermined amount based on the field capacity.

A concentration of 0.3% manganese sulfate was used for priming the seeds of both sesame cultivars. Thus, the seeds of both sesame cultivars were immersed in distilled water (as the control) and

manganese sulfate solution at a ratio of 1:5 (one gram of seed in 5 milliliters of solution) for seven hours. Afterward, the seeds were placed on filter paper to achieve the initial moisture level. Foliar spraying of MnSO₄•7H₂O was applied twice: 48 hours before exposure to water deficiency and 7 days after the onset of water stress. For foliar spraying of manganese sulfate, manganese sulfate was used at a concentration of 0.3% for the Dashtestan2 and Naz Takshakheh cultivars. Foliar spraying was done using a 2-liter sprayer. Foliar spraying was conducted in a way that covered the entire leaf surface of the plant. Control treatments were sprayed using distilled water. Leaf sampling was conducted 48 hours after the second foliar spray at 10 AM, the third fresh leaf of each plant was harvested and stored at -80°C for measurement of biochemical indicators, including activities of CAT, POX, APX, as well as the concentrations of proline, soluble sugar, and soluble protein. To evaluate seed yield, sampling was performed at the maturity stage.

CAT enzyme activity was measured using the method of Aebi (1984). In order to measure the CAT enzyme activity, first, the spectrophotometer was set to a wavelength of 240 nm. Then the microtubes containing the extracted enzyme were taken out of the refrigerator and placed on ice. Then 3000 µl of 100 mM phosphate buffer and 40 µl of 30 mM hydrogen peroxide were poured into a quartz cell. The cell containing these two substances was placed in the spectrophotometer and used to calibrate the device. Next, 100 µl of enzyme extract was added to it and after five minutes of starting the reaction at room temperature, absorption was performed using a spectrophotometer at a wavelength of 240 nm. The method of MacAdam et al. (1992) was used to measure POX enzyme activity. To measure the activity of the POX enzyme, first, the spectrophotometer was set to a wavelength of 470 nm. Then the microtubes containing the extracted enzyme were taken out of the refrigerator and placed on ice. Then, 3 ml of 100 mM phosphate buffer, 50 µl of 200 mM guaiacol, and 40 µl of 30 mM hydrogen peroxide were mixed. Then 100 µl of enzyme extract was added to each sample and five minutes after

the start of the reaction at laboratory temperature, absorption was performed at a wavelength of 470 nm using a spectrophotometer. The activity of the POX enzyme was expressed as enzyme units per gram of fresh weight (FW). APX enzyme activity was measured by the method of Nakano and Asada (1987). The reaction mixture consisted of 1500 µl of 50 mM phosphate buffer (pH 7.8), 400 µl of 0.1 mM hydrogen peroxide, 400 µl of 30% ascorbic acid, 600 µl of 0.1 mM EDTA, and 100 µl of enzyme extract. APX enzyme activity was calculated based on the rate of hydrogen peroxide decomposition per minute per mg of protein at a wavelength of 290 nm and using an extinction coefficient of $2.8 \text{ mM}^{-1}\text{cm}^{-1}$. In order to calculate seed yield, all plants in each pot (5 plants) were harvested 100 and 105 days after planting in the

Naz Takshakheh and Dashtestan2 cultivars, respectively, and after separating the seed from the plants, the seed yield per plant was calculated. Before conducting the statistical analysis of the data and analysis of variance, normal distribution tests for the data and errors were performed using the Kolmogorov, Immonrov, and Levin methods with SPSS software. The comparison of main effects means was conducted using Duncan's multiple range method at a probability level of 5% with MSTATC software.

3. Results and discussion

The analysis of variance showed that the individual effects of cultivar, spray, Seed Priming and water deficiency on the examined traits were statistically significant ($p \leq 0.01$) (Table 3).

Table 3. Analysis of variance of traits of two sesame plant cultivars in response to manganese sulfate application methods under water deficiency

S.O.V.	df	Mean Squares						
		Seed yield per plant	Proline	Soluble sugar content	Soluble protein content	CAT activity	POX activity	APX activity
Replication	2	0.00015 ^{ns}	0.00041 ^{ns}	0.00011 ^{ns}	0.0016 ^{ns}	0.00066 ^{ns}	0.00002 ^{ns}	0.000031 ^{ns}
Cultivar (C)	1	0.195 ^{**}	0.00097 [*]	0.00087 ^{**}	0.471 ^{**}	0.0224 ^{**}	0.0074 [*]	0.144 ^{**}
Spray (S)	1	0.53 ^{**}	0.156 ^{**}	0.512 ^{**}	0.4036 ^{**}	0.278 ^{**}	0.307 ^{**}	0.45 ^{**}
Priming (P)	1	0.127 ^{**}	0.046 ^{**}	0.168 ^{**}	0.1092 ^{**}	0.15 ^{**}	0.115 ^{**}	0.18 ^{**}
Water deficiency (W)	2	0.634 ^{**}	0.452 ^{**}	0.15 ^{**}	0.4368 ^{**}	0.608 ^{**}	0.165 ^{**}	0.496 ^{**}
C × S	1	0.052 ^{**}	0.000015 ^{ns}	0.00056 [*]	0.00044 ^{ns}	0.0004 ^{ns}	0.0012 ^{**}	0.0095 ^{ns}
C × P	1	0.008 ^{**}	0.0000023 ^{ns}	0.0014 ^{**}	0.007 [*]	0.00015 ^{ns}	0.00035 ^{**}	0.0034 ^{ns}
C × W	2	0.0014 ^{**}	0.00000019 ^{ns}	0.0014 ^{**}	0.0366 ^{**}	0.000006 ^{ns}	0.00023 ^{**}	0.00074 ^{**}
S × P	1	0.0052 ^{**}	0.000044 ^{ns}	0.043 ^{**}	0.000015 ^{ns}	0.0098 ^{**}	0.000034 ^{ns}	0.0295 ^{**}
S × W	2	0.00036 ^{ns}	0.0067 ^{**}	0.00008 ^{ns}	0.0148 ^{**}	0.0101 ^{**}	0.0015 ^{**}	0.0737 ^{**}
P × W	2	0.005 ^{**}	0.0017 ^{**}	0.0002 ^{ns}	0.0156 ^{**}	0.0054 ^{**}	0.00091 ^{**}	0.0299 ^{**}
C × S × P	1	0.0042 ^{**}	0.0000023 ^{ns}	0.011 ^{**}	0.00029 ^{ns}	0.000063 ^{ns}	0.00034 ^{**}	0.0018 ^{ns}
C × S × W	2	0.0006 [*]	0.0000083 ^{ns}	0.00009 ^{ns}	0.00069 ^{ns}	0.00047 ^{ns}	0.0007 ^{**}	0.0027 ^{ns}
C × P × W	2	0.0024 ^{**}	0.000014 ^{ns}	0.0002 ^{ns}	0.00064 ^{ns}	0.0025 ^{ns}	0.00034 ^{**}	0.0009 ^{ns}
S × P × W	2	0.0045 ^{**}	0.000056 ^{ns}	0.0005 ^{**}	0.002 ^{ns}	0.0182 ^{**}	0.00084 ^{**}	0.0024 ^{ns}
C × S × P × W	2	0.0047 ^{**}	0.000025 ^{ns}	0.00012 ^{**}	0.0015 ^{ns}	0.0027 ^{ns}	0.00067 ^{**}	0.00051 ^{ns}
Error	46	0.0001	0.00021	0.00008	0.0011	0.00088	0.000038	0.0035
C.V. (%)		2.39	3.76	1.51	8.28	4.23	1.15	18.31

3.1. Seed yield per plant

The mean comparison indicated that the seed yield of both sesame cultivars was adversely affected by rising water deficiency levels. The reduction in seed yield can be attributed to the negative effect of drought stress on the seed number per plant and the weight of 1000 seeds, which was linked to a shortened duration of the seed filling process (Alizadeh et al., 2010). The seed yield of Dashtestan2 and Naz Takshakheh cultivars decreased from 0.568 grams and 0.526 grams, respectively, under well-watered conditions (75% FC) to 0.241 grams and 0.218 grams per plant under severe drought conditions (25% of FC) (Fig. 1).




Figure 1. The effect of different manganese sulfate applications on seed yield of two sesame cultivars under water-deficient conditions. Different letters indicate significant differences ($p < 0.05$, Duncan's test).

The most noticeable impact of water deficiency on plants is the decrease in leaf growth and development, caused by slowed growth and cell division. This reduction subsequently leads to lower dry matter accumulation and crop yield. Additionally, a study showed that different soybean genotypes experienced a significant reduction in seed yield under water-deficient conditions (Soliman *et al.*, 2025). At all levels of water deficiency, seed yield was higher in the Dashtestan2 cultivar than in the Naz Takshakheh cultivar. Priming seeds with nutrients enhances plant establishment, accelerates phenological development, boosts yield, and improves the micronutrient concentration in crops. Additionally, in many instances, priming seeds with micronutrients can decrease the requirement for these nutrients and reduce dependence on fertilizers. Seed priming is a straightforward and cost-effective technique for nutrient application (Farooq *et al.*, 2012).

The findings indicated that utilizing manganese sulfate through a combination of seed priming and solution spraying proved to be more effective in mitigating the impact of water scarcity on seed yield than employing each method independently. The highest seed yield, $0.879 \text{ g plant}^{-1}$, was obtained from the Dashtestan2 cultivar under well-watered conditions (75% FC) using the seed priming and solution spraying treatment. In contrast, the lowest seed yield, 0.218 g/plant , was recorded for the Naz Takshakheh cultivar under severe water deficiency conditions (25% FC) without any treatment (control) (Fig. 1). The results of this study showed that the application of manganese sulfate in the form of seed priming, foliar spraying, and seed priming + solution spraying significantly and remarkably increased the seed yield per plant of both sesame cultivars at all levels of drought stress. The enhancement in seed yield from the application of iron, zinc, and manganese can be linked to these elements' ability to boost the activity of antioxidant enzymes, including SOD, CAT, and POX, which help mitigate ROS (Waraich *et al.*, 2012).

Furthermore, another study found that foliar spraying of micronutrients led to improved wheat yield and its components (Shoormij *et al.*, 2022). Other research results showed that yield and yield components in different canola cultivars decreased under late-season drought stress (Khayat Moghadam *et al.*, 2021). Under water-deficient stress conditions, the

application of zinc and manganese through foliar treatments on canola (Khodabin *et al.*, 2022) and safflower (Movahhedy-Dehnavy *et al.*, 2009) resulted in increased seed yield.

3.2. Proline content

The research results showed that as the levels of water deficiency escalated, there was a corresponding increase in the proline concentration found in sesame leaves. Specifically, proline levels increased from 0.224 mg/g of leaf FW under well-watered conditions (75% of FC) to 0.473 mg/g of leaf FW at the severe water deficiency level (25% of FC) (Fig. 2). Under water deficiency stress conditions, plants accumulate organic osmolytes to maintain osmotic balance (Blum, 2017). A high concentration of osmoprotectants, including proline, sugars, polyamines, and glycine betaine, is generally associated with increased biosynthesis and improved stress tolerance in plants (Kaur *et al.*, 2017). Proline serves a dual function: at low to moderate concentrations, it has beneficial effects and plays a role in signaling, while at higher concentrations, it can be toxic to living cells. Beyond osmotic regulation, osmoprotectants also perform various other functions, including scavenging free radicals (Szabados and Savouré, 2010), inducing the expression of osmotic stress-related genes, and stabilizing cell membranes and proteins (Theocharis *et al.*, 2012). Many studies have indicated an increase in proline levels due to water deficiency stress in various plants, including wheat (Kamruzzaman *et al.*, 2023), rapeseed (Farahani *et al.*, 2020), sorghum (Amoah and Antwi-Berko, 2020), and chickpea (Seifikalhor *et al.*, 2022).

Comparisons of the mean data indicated that the foliar application of manganese sulfate increased proline levels in sesame plants at all levels of water deficiency. Proline concentration increased by 0.085 , 0.065 , and 0.13 mg/g of leaf FW at 75%, 50%, and 25% of FC, respectively, compared to the control treatment (no foliar application) (Fig. 2). Overall, these results indicate that the foliar application of manganese sulfate enhances drought tolerance. Several studies have linked the increase in proline levels from foliar applications of zinc and manganese to the activation of protein-degrading enzymes, which convert proteins into the amino acid proline (Babaeian *et al.*, 2011; Waraich *et al.*, 2012).




Figure 2. The effect of foliar spraying with manganese sulfate on proline content in sesame leaves under water deficiency conditions. Different letters indicate significant differences ($p < 0.05$, Duncan's test).

Proline concentration in sesame leaves increased with seed priming under all levels of water deficiency. Priming sesame seeds with manganese sulfate led to a more significant rise in proline levels at all three levels of water deficiency compared to the control condition without priming. Specifically, seed priming led to increases in proline of 0.048, 0.036, and 0.069 mg g⁻¹ of FW at 75%, 50%, and 25% of FC, respectively, compared to the control (Fig. 3). Proline accumulation under water deficit conditions functions as an osmolyte and enhances the ability to repair damage (Kumar et al., 2011; Anjum et al., 2011). Water deficit stress results in increased proline accumulation in wheat (Sawhney and Singh, 2002) and soybean (Krüger, 2002). Additionally, manganese application has been reported to increase proline content in wheat leaves subjected to drought stress (Ahmad et al., 2019).




Figure 3. The effect of seed priming with manganese sulfate on leaf proline content under water deficiency conditions. Different letters indicate significant differences ($p < 0.05$, Duncan's test).

3.3. Soluble sugar

Increasing water deficiency intensity elevated the soluble sugar content in both sesame cultivars. In the Dashtestan2 and Naz Takshakheh cultivars, the soluble

sugar content increased from 0.406 and 0.374 mg g⁻¹ leaf dry weight (DW) under well-watered conditions to 0.478 and 0.448 mg g⁻¹ leaf DW at mild water deficiency (50% FC), respectively, reaching 0.568 and 0.543 mg g⁻¹ leaf DW under severe water deficiency (25% FC) conditions (Fig. 4). Other studies have found that drought stress conditions increase the levels of soluble sugars in various plants, including corn (Zi et al., 2022), rice (Wang et al., 2019), wheat (Johari-Pireivatlou, 2010), rapeseed (Moradshahi et al., 2004), and soybean (Du et al., 2020). The activity of the alpha-amylase enzyme seems to increase under water deficiency stress, resulting in a higher concentration of soluble sugars due to starch hydrolysis (Setter et al., 2001). During periods of water deficiency, the carbohydrate content in the plant rises, mainly serving functions in osmotic regulation, protection, and carbon storage (Cechin et al., 2006). These sugars act as a defense mechanism during water deficiency stress, helping to maintain elevated turgor pressure within the cells (Kishor et al., 2005).

Increasing water deficiency stress and employing various methods of manganese sulfate application—such as seed priming, foliar spraying, and a combination of both—resulted in a higher concentration of soluble sugars in the leaves of both sesame cultivars across all three levels of water deficiency stress compared to the control treatment (which did not receive manganese sulfate treatment). The Dashtestan2 cultivar demonstrated the highest concentration of soluble sugar, measuring 0.838 mg g⁻¹ dry leaf weight, when treated with manganese sulfate via foliar spraying and priming at severe water deficiency (25% FC). In contrast, the Naz Takshakheh cultivar exhibited the lowest concentration of soluble sugar, at 0.374 mg g⁻¹ dry leaf weight, under well-watered conditions with manganese sulfate treatment (Fig. 4). Foliar application of 400 mg L⁻¹ manganese sulfate enhanced the levels of total carbohydrates, total protein, and antioxidant status in two cultivars of grapevine (*Vitis vinifera* cv. Rotabi, and Thompson seedless) subjected to water deficiency stress (Ghorbani et al., 2019). The increase in carbohydrate production in some plants treated with manganese sulfate may be linked to their higher chlorophyll content. Carbohydrate production is influenced by both the amount of chlorophyll and its efficiency (Lidon et al., 2004; Millaleo et al., 2010).




Figure 4. The effect of different methods of manganese sulfate application on the soluble sugar content of two sesame cultivars under water-deficient stress conditions. Different letters indicate significant differences ($p < 0.05$, Duncan's test).

3.4. Soluble protein

The mean comparison of data revealed a difference in soluble protein content between the two sesame cultivars. In the control treatment, the concentration of leaf soluble protein was $0.448 \text{ mg g}^{-1} \text{ FW}$ for the Dashtestan2 cultivar, while it was $0.305 \text{ mg g}^{-1} \text{ FW}$ for the Naz Takshakheh cultivar (Fig. 5). The experiment results indicated that seed priming with manganese sulfate enhanced the concentration of leaf soluble protein in both the Dashtestan2 and Naz Takshakheh cultivars. Specifically, the concentrations increased by 0.097 mg g^{-1} of leaf FW for Dashtestan2 and 0.059 mg g^{-1} for Naz Takshakheh, compared to the control treatment. These findings align with previous studies on wheat (Ahmad et al., 2019).




Figure 5. The effect of seed priming on the soluble protein content of two sesame cultivars. Different letters indicate significant differences ($p < 0.05$, Duncan's test).

As water deficiency stress levels increased, the soluble protein content in both sesame cultivars decreased. In the Dashtestan2 and Naz Takshakheh cultivars, soluble protein levels dropped from 0.601

and 0.527 mg g^{-1} leaf FW under well-watered conditions to 0.411 and 0.188 mg g^{-1} leaf FW, respectively, under severe water deficiency (25% FC) conditions (Fig. 6). The results indicated that at varying levels of water deficiency stress, the soluble protein content of the Dashtestan2 cultivar was consistently higher than that of the Naz Takshakheh cultivar, with values of 0.074 , 0.188 , and 0.223 mg g^{-1} of leaf FW, respectively (Fig. 6). Additionally, a study on wheat revealed that water deficiency stress conditions lead to a decrease in the protein content of wheat leaves (Ahmad et al., 2019).




Figure 6. The effect of water deficiency stress on the soluble protein content of two sesame cultivars. Different letters indicate significant differences ($p < 0.05$, Duncan's test).

The results demonstrate the beneficial effect of manganese sulfate foliar application on soluble protein levels under varying water deficiency conditions. As water deficiency stress intensified, the concentration of soluble protein in the leaves decreased. However, the application of manganese sulfate as a foliar treatment enhanced the soluble protein concentration at three levels of water deficiency stress, resulting in increases of 0.207 , 0.126 , and 0.116 mg g^{-1} of leaf FW, respectively, compared to the control treatment (without foliar application) (Fig. 7). Fathi Amirkhiz et al. (2015) reported that the lack of manganese foliar application led to a decrease in protein content under water deficiency stress conditions. Manganese is effective in enhancing photosynthetic carbon conversion and drought tolerance by promoting the synthesis of soluble proteins and facilitating photosynthesis (Jafardokht et al., 2015). Foliar application of manganese sulfate can increase the protein content of grape leaves under water deficiency stress conditions (Lidon et al., 2004; Millaleo et al., 2010).




Figure 7. The effect of foliar spraying of plants with manganese sulfate on the amount of leaf soluble protein under water deficiency stress conditions. Different letters indicate significant differences ($p < 0.05$, Duncan's test).

Seed priming reduced the negative impact of water deficiency stress on the concentration of soluble leaf protein. Specifically, under three levels of water deficiency stress, soluble sesame leaf protein increased by 0.137, 0.049, and 0.048 mg g^{-1} of leaf FW, respectively, due to seed priming with manganese sulfate compared to the control treatment (without priming) (Fig. 8).




Figure 8. The effect of seed priming with manganese sulfate on leaf soluble protein content under drought stress conditions. Different letters indicate significant differences ($p < 0.05$, Duncan's test).

3.5. CAT enzyme activity

The results indicated that applying manganese sulfate through seed priming and foliar spraying enhanced CAT activity. Specifically, seed priming increased CAT activity by approximately 0.115 units, while foliar spraying resulted in a greater increase of about 0.148 units compared to the control treatment (Fig. 9). Manganese, an essential micronutrient, plays a crucial role in minimizing the production of oxygen-free radicals in plants caused by environmental stresses. It achieves this directly by enhancing the

composition and activity of antioxidant enzymes, and indirectly by increasing the rates of photosynthesis and nitrogen metabolism (Waraich et al., 2012).




Figure 9. The interaction effect of seed priming and manganese sulfate foliar application on CAT activity. Different letters indicate significant differences ($p < 0.05$, Duncan's test).

The results indicated that foliar application of manganese sulfate had a significantly greater impact on CAT activity than the seed priming method. Furthermore, CAT enzyme activity was higher in plants grown from primed seeds that also received foliar spraying with manganese sulfate, compared to those that underwent only seed priming or foliar spray treatment alone. As water deficiency stress levels increased, CAT enzyme activity also increased. The combined application of manganese sulfate through priming and foliar spraying further enhanced CAT activity under conditions of water deficiency stress. In the seed priming method, the activity of CAT enzyme increased by 0.208, 0.048, and 0.09 units compared to the control conditions (without seed priming and foliar spraying) at 75%, 50%, and 25% of field capacity, respectively (Fig. 10).




Figure 10. The interaction effect of seed priming and manganese sulfate foliar application on CAT activity. Different letters indicate significant differences ($p < 0.05$, Duncan's test).

In the foliar spraying method, the increases were 0.247, 0.095, and 0.102 units, compared to the control conditions (without seed priming and foliar spraying). Additionally, the level of CAT enzyme activity increased more in the combined application method of seed priming and foliar spraying with manganese sulfate than in their separate application (Fig. 10). CAT is a heme-containing enzyme that catalyzes the conversion of hydrogen peroxide (H_2O_2) into water (H_2O) and oxygen (O_2) (Anjum et al., 2016). During water deficiency stress conditions, the activity of CAT can vary; it may increase, decrease, or stay the same (Mishra and Panda, 2017). Increased activity of the CAT enzyme during water deficiency stress conditions helps eliminate H_2O_2 , which in turn reduces cell damage and enhances the oxidative capacity of plants (Piri et al., 2019). Antioxidant enzymes, such as CAT and SOD in safflower (Zafari et al., 2020) and CAT and APX in castor (de Araújo Silva et al., 2016), increased under water deficiency stress.

3.6. POX enzyme activity

The results indicated a difference in POX enzyme activity between the two sesame cultivars under all levels of water deficiency stress and manganese sulfate application methods. Specifically, the Dashtestan2 cultivar exhibited higher POX enzyme activity compared to the Naz Takshakheh cultivar. Various methods of applying manganese sulfate enhanced POX enzyme activity under all levels of water deficiency stress. Notably, the combined approach of seed priming and foliar spraying with manganese sulfate resulted in higher POX activity compared to each method applied separately. As water deficiency stress increased, POX enzyme activity rose in both sesame cultivars (Fig. 11). Similar findings have been reported in various plants, including maize (Shafiq et al., 2021), faba bean (Siddiqui et al., 2015), and mung bean (Abid et al., 2017) under water deficiency stress conditions. Many researchers believe that the proper functioning of various enzymes, such as CAT, POX, and SOD, helps reduce the disruption of cellular stability caused by ROS (Hussain et al., 2016; Ashraf, 2009; Moradi and Piri, 2018).

3.7. APX enzyme activity

Increasing water deficiency stress levels correlated with elevated activity of the APX enzyme in both

sesame cultivars. However, there was a notable difference in enzyme activity between the two cultivars at all water deficiency stress levels. Specifically, in the Dashtestan2 cultivar, the APX activity at 75%, 50%, and 25% of FC was 0.081, 0.086, and 0.102 units higher than that of the Naz Takshakheh cultivar, respectively (Fig. 12). Studies on safflower (Khosrowshahi et al., 2020) and milk thistle (ElSayed et al., 2019) have indicated that water deficiency stress results in an increase in the activity of the APX enzyme. Similarly, research involving two canola cultivars—one sensitive and one tolerant—demonstrated that both cultivars exhibited increased APX enzyme activity under water deficiency stress conditions (Seyed Ebrahimi et al., 2018).




Figure 11. The effect of different methods of manganese sulfate application on the POX activity of two sesame cultivars under water deficiency stress conditions. Different letters indicate significant differences ($p < 0.05$, Duncan's test).




Figure 12. The effect of water deficiency stress on the activity of APX enzyme in two sesame cultivars. Different letters indicate significant differences ($p < 0.05$, Duncan's test).

APX enzyme activity increased due to seed priming and the foliar application of manganese sulfate. Specifically, enzyme activity rose by 0.06, 0.138, and 0.258 units after seed priming, foliar spraying, and

combined priming with solution spraying compared to the control treatment, respectively (Fig. 13). Similarly, a study by Jalili and Karimi (2022) on grapes found that foliar spraying with manganese also enhances POX enzyme activity.




Figure 13. The interaction effect of seed priming and foliar spraying with manganese sulfate on the activity of the APX enzyme. Different letters indicate significant differences ($p < 0.05$, Duncan's test).

As the level of water deficiency stress increased, the activity of APX increased from 0.151 units in the well watered condition (75% of FC) to 0.339 units in the severe water deficiency condition (25% FC). Foliar spraying of the plant with manganese sulfate led to an increase in the activity of APX at all levels of water deficiency stress compared to the control treatment. The activity of the APX enzyme increased by 0.097, 0.09, and 0.286 units compared to the control, with foliar spraying applied at 75, 50, and 25% of FC, respectively (Fig. 14).




Figure 14. The effect of manganese sulfate foliar application on APX enzyme activity under water deficiency stress conditions. Different letters indicate significant differences ($p < 0.05$, Duncan's test).

Compared to non-primed seeds, manganese sulfate priming boosted APX enzymatic function regardless of drought severity level. Specifically, seed priming at well watered (75% FC) increased enzyme activity by

0.077 units, at mild water deficiency (50% FC) by 0.044 units, and at severe water deficiency (25% FC) by 0.179 units compared to the control treatment (Fig. 15).




Figure 15. The effect of seed priming with manganese sulfate on APX enzyme activity under water-deficient stress conditions. Different letters indicate significant differences ($p < 0.05$, Duncan's test).

4. Conclusion

This study found that water deficiency stress reduced both the seed yield and the soluble leaf protein content of the two sesame cultivars. Drought stress induced a dual response: osmotic adjustment via proline and sugar accumulation, coupled with enhanced antioxidant activity, such as CAT, POX, and APX activity. This study demonstrates that the combined seed priming and foliar application of MnSO_4 significantly outperformed either method applied individually, establishing a novel agronomic strategy for drought mitigation in sesame. Notably, cultivar-specific genetic variations in stress adaptation were identified, with Dashtestan2 exhibiting superior antioxidant defense activation and yield stability under water deficit conditions compared to Naz Takshakheh. Furthermore, the mechanistic linkage between MnSO_4 -induced proline accumulation and enhanced antioxidant enzyme activity provides critical insights into micronutrient-mediated drought resilience, advancing our understanding of plant stress physiology. Future studies should explore the molecular basis of MnSO_4 -induced drought tolerance in sesame, including transcriptomic analyses of stress-responsive pathways, while also optimizing field-level application protocols for diverse cultivars and environments. Additionally, investigating synergistic effects with other micronutrients or bio-stimulants could further enhance drought resilience strategies for sustainable sesame production under climate change.

Conflict of interests

All authors declare no conflict of interest.

Ethics approval and consent to participate

No humans or animals were used in the present research. The authors have adhered to ethical standards, including avoiding plagiarism, data fabrication, and double publication.

Consent for publications

All authors read and approved the final manuscript for publication.

Availability of data and material

All the data are embedded in the manuscript.

Authors' contributions

All authors had an equal role in study design, work, statistical analysis and manuscript writing.

Informed consent

The authors declare not to use any patients in this research.

Funding/Support

This study was supported by Ilam University, Ilam, Iran.

Acknowledgement

This article was achieved based on the material and equipment of Department of Agronomy and Crop Breeding, Faculty of Agriculture, Ilam University, that the authors thanks it.

References

- Abid G., M'hamdi M., Mingeot D., Aouida M., Aroua I., Muhovski Y., Sassi K., Souissi F., Mannai K., Jebara M. 2017. Effect of drought stress on chlorophyll fluorescence, antioxidant enzyme activities and gene expression patterns in faba bean (*Vicia faba* L.). Archives of Agronomy and Soil Science 63(4): 536-552. <https://doi.org/10.1080/03650340.2016.1224857>
- Aebi H. 1984. Catalase in vitro. In Methods in Enzymology (105: pp. 121-126). Academic Press. [https://doi.org/10.1016/S0076-6879\(84\)05016-3](https://doi.org/10.1016/S0076-6879(84)05016-3)
- Ahmad I., Kamran M., Yang X., Meng X., Ali S., Ahmad S., Zhang X., Bilegjargal B., Ahmad B., Liu T., Cai T. 2019. Effects of applying uniconazole alone or combined with manganese on the photosynthetic efficiency, antioxidant defense system, and yield in wheat in semiarid regions. Agricultural Water Management 216: 400-414. <https://doi.org/10.1016/j.agwat.2019.02.025>
- Alizadeh A., Khoshkhui M., Javidnia K., Firuzi O., Tafazoli E., Khalighi A. 2010. Effects of fertilizer on yield, essential oil composition, total phenolic content and antioxidant activity in *Satureja hortensis* L.(Lamiaceae) cultivated in Iran. Journal of Medicinal Plants Research 4(1): 33-40. <https://doi.org/10.5897/JMPR09.361>
- Amoah J.N., Antwi-Berko D. 2020. Comparative physiological, biochemical and transcript response to drought in sorghum genotypes. Biotechnology Journal International 24(3): 1-14. <https://doi.org/10.9734/BJI/2020/v24i330102>
- Amritha M.S., Sridharan K., Puthur J.T., Parkash Dhankher O.M. 2021. Priming with nanoscale materials for boosting abiotic stress tolerance in crop plants. Journal of Agricultural and Food Chemistry 69(35): 10017-10035. <https://doi.org/10.1021/acs.jafc.1c03673>
- Anjum N.A., Sharma P., Gill S.S., Hasanuzzaman M., Khan E.A., Kachhap K., Mohamed A.A., Thangavel P., Devi G.D., Vasudhevan P., Sofo A. 2016. Catalase and ascorbate peroxidase—representative H₂O₂-detoxifying heme enzymes in plants. Environmental Science and Pollution Research 23(19): 19002-19029. <https://doi.org/10.1007/s11356-016-7309-6>
- Anjum S.A., Xie X., Wang L.C., Saleem M.F., Man C., Lei W. 2011. Morphological, physiological and biochemical responses of plants to drought stress. African Journal of Agricultural Research 6(9): 2026-2032. <https://doi.org/10.5897/AJAR10.02>
- Ashraf M.J. 2009. Biotechnological approach of improving plant salt tolerance using antioxidants as markers. Biotechnology Advances 27(1): 84-93. <https://doi.org/10.1016/j.biotechadv.2008.09.003>
- Babaeian M., Tavassoli A., Ghanbari A., Esmailian Y., Fahimifard M. 2011. Effects of foliar micronutrient application on osmotic adjustments, grain yield and yield components in sunflower (Alstar cultivar) under water stress at three stages. African Journal of Agricultural Research 6(5): 1204-1208. <https://doi.org/10.5897/AJAR10.928>
- Balouchi H., Soltani Khankahdani V., Moradi A., Gholamhoseini M., Piri R., Heydari S.Z., Dedicova B. 2023. Seed fatty acid changes germination response to temperature and water potentials in six sesame (*Sesamum indicum* L.) cultivars: estimating the cardinal temperatures. Agriculture 13(10): 1936. <https://doi.org/10.3390/agriculture13101936>
- Blum A. 2017. Osmotic adjustment is a prime drought stress adaptive engine in support of plant production. Plant, Cell & Environment 40(1): 4-10. <https://doi.org/10.1111/pce.12800>
- Cechin I., Rossi S.C., Oliveira V.C., Fumis T.D. 2006. Photosynthetic responses and proline content of mature and young leaves of sunflower plants under water deficit. Photosynthetica 44(1): 143-146. <https://doi.org/10.1007/s11099-005-0171-2>
- Chen J., Yang L., Yan X., Liu Y., Wang R., Fan T., Ren Y., Tang X., Xiao F., Liu Y., Cao S. 2016. Zinc-finger transcription factor ZAT6 positively regulates cadmium tolerance through the glutathione-dependent pathway in Arabidopsis. Plant Physiology 171(1): 707-719. <https://doi.org/10.1104/pp.15.01882>
- Dawood M.F., Abeer A.H., Aldaby E.E. 2019. Titanium dioxide nanoparticles model growth kinetic traits of some wheat

- cultivars under different water regimes. *Plant Physiology Reports* 24(1): 129-140. <https://doi.org/10.1007/s40502-019-0437-5>
- de Araújo Silva M.M., Willadino L., dos Santos D.Y., Oliveira A.F., Camara T.R. 2016. Response of *Ricinus communis* L. to in vitro water stress induced by polyethylene glycol. *Plant Growth Regulation* 78(2): 195-204. <https://doi.org/10.1007/s10725-015-0085-3>
- Diya A., Beena R., Jayalekshmy V.G. 2024. Physiological, biochemical and molecular mechanisms of seed priming: a review. *Legume Research: An International Journal* 47(2): 159-166. <https://doi.org/10.18805/LR-4638>
- Du Y., Zhao Q., Chen L., Yao X., Zhang W., Zhang B., Xie F. 2020. Effect of drought stress on sugar metabolism in leaves and roots of soybean seedlings. *Plant Physiology and Biochemistry* 146: 1-12. <https://doi.org/10.1016/j.plaphy.2019.11.003>
- ElSayed A.I., El-Hamahmy M.A., Rafudeen M.S., Mohamed A.H., Omar A.A. 2019. The impact of drought stress on antioxidant responses and accumulation of flavonolignans in milk thistle (*Silybum marianum* (L.) Gaertn). *Plants* 8(12): 611. <https://doi.org/10.3390/plants8120611>
- Farahani S., Shahsavari N., Mohammadi Arasteh M. 2020. Effect of potassium sulfate on the physiological characteristics of canola cultivars in late season drought stress conditions. *Journal of Plant Nutrition* 43(9): 1217-1228. <https://doi.org/10.1080/01904167.2020.1727507>
- Farooq M., Usman M., Nadeem F., Rehman H.U., Wahid A., Basra S.M., Siddique K.H. 2019. Seed priming in field crops: potential benefits, adoption and challenges. *Crop & Pasture Science* 70(9): 731-771. <https://doi.org/10.1071/CP18604>
- Farooq M., Wahid A., Kadambot H.M.S. 2012. Micronutrient application through seed treatments, a review. *Journal of Soil Science and Plant Nutrition* 12: 125-142. <https://doi.org/10.4067/S0718-95162012000100011>
- Fathi Amirkhiz K., Amini Dehaghi M., Heshmati S. 2015. Study the effect of iron chelate on chlorophyll content, photochemical efficiency and some biochemical traits in safflower under deficit irrigation condition. *Iranian Journal of Field Crop Science* 46(1): 137-145. (In Farsi). <https://doi.org/10.22059/ijfcs.2015.54053>
- Ghorbani P., Eshghi S., Ershadi A., Shekafandeh A., Razzaghi F. 2019. The possible role of foliar application of manganese sulfate on mitigating adverse effects of water stress in grapevine. *Communications in Soil Science and Plant Analysis* 50(13): 1550-1562. <https://doi.org/10.1080/00103624.2019.1626873>
- Ghosh S., Bera K., Dutta P., Sadhukhan S. 2023. Seed priming and seedling pre-treatment in regulating secondary metabolism for stress tolerance. In *Biology and Biotechnology of Environmental Stress Tolerance in Plants* (pp. 263-304). Apple Academic Press.
- Gray S.B., Brady S.M. 2016. Plant developmental responses to climate change. *Developmental Biology* 419(1): 64-77. <https://doi.org/10.1016/j.ydbio.2016.07.023>
- Hamarash H., Hamad R., Rasul A. 2022. Meteorological drought in semi-arid regions: A case study of Iran. *Journal of Arid Land* 14(11): 1212-1233. <https://doi.org/10.1007/s40333-022-0106-9>
- Hasanuzzaman M., Bhuyan M.B., Anee T.I., Parvin K., Nahar K., Mahmud J.A., Fujita M. 2019. Regulation of ascorbate-glutathione pathway in mitigating oxidative damage in plants under abiotic stress. *Antioxidants* 8(9): 384. <https://doi.org/10.3390/antiox8090384>
- Hernandez-Leon S.G., Valenzuela-Soto E.M. 2023. Glycine betaine is a phytohormone-like plant growth and development regulator under stress conditions. *Journal of Plant Growth Regulation* 42(8): 5029-5040. <https://doi.org/10.1007/s00344-022-10855-3>
- Hussain M., Ul-Allah S., Farooq S. 2023. Sesame (*Sesamum indicum* L.). In *Neglected and Underutilized Crops* (pp. 733-755). Academic Press. <https://doi.org/10.1016/B978-0-323-90537-4.00026-0>
- Hussain S., Khan F., Cao W., Wu L., Geng M. 2016. Seed priming alters the production and detoxification of reactive oxygen intermediates in rice seedlings grown under sub-optimal temperature and nutrient supply. *Frontiers in Plant Science* 7: 188539. <https://doi.org/10.3389/fpls.2016.00439>
- Jafardokht R., Mosavi Nik S.M., Mehraban A., Basiri M. 2015. Effect of water stress and foliar micronutrient application on physiological characteristics and nutrient uptake in mung bean. *Electronic Journal of Crop Production* 8(1): 121-141. (In Farsi). <https://dor.isc.ac/dor/20.1001.1.2008739.1394.8.1.7.7>
- Jalili M., Karimi R. 2022. The effect of combined application of nitrogen, zinc and manganese on yield and quality characteristics of grape fruit. *Plant Biological Sciences* 13(4): 33-56. (In Farsi). <https://doi.org/10.22108/ijpb.2022.133403.1284>
- Jogawat A. 2019. Osmolytes and their role in abiotic stress tolerance in plants. *Molecular Plant Abiotic Stress: Biology and Biotechnology* (pp. 91-104). <https://doi.org/10.1002/9781119463665.ch5>
- Johari-Pireivatlou M. 2010. Effect of soil water stress on yield and proline content of four wheat lines. *African Journal of Biotechnology* 9(1): 36-40. <https://www.ajol.info/index.php/ajb/article/view/77766>
- Kamruzzaman M., Shrestha A., Siddiqui M.N., Oyiga B.C., Ballvora A., Léon J., Naz A.A. 2023. Genetic mapping of candidate loci for water-deficit stress-induced proline accumulation in bread wheat (*Triticum aestivum*). *Plant Breeding* 142(4): 449-462. <https://doi.org/10.1111/pbr.13096>
- Kar R.K. 2011. Plant responses to water stress: role of reactive oxygen species. *Plant Signaling & Behavior* 6(11): 1741-1745. <https://doi.org/10.4161/psb.6.11.17729>
- Kaur D., Grewal S.K., Kaur J., Singh S. 2017. Differential proline metabolism in vegetative and reproductive tissues determine drought tolerance in chickpea. *Biologia Plantarum* 61(2): 359-366. <https://doi.org/10.1007/s10535-016-0695-2>
- Kermani S.G., Saeidi G., Sabzalian M.R., Gianinetti A. 2019. Drought stress influenced sesamin and sesamol content and polyphenolic components in sesame (*Sesamum indicum* L.) populations with contrasting seed coat colors. *Food Chemistry* 289: 360-368. <https://doi.org/10.1016/j.foodchem.2019.03.004>
- Khayat Moghadam M.S., Gholami A., Shirani Rad A.H., BaradaranFiroozabadi M., Abbasdokht H. 2021. The effect of potassium silicate and late-season drought stress on the physiological characters of canola. *Journal of Crops*

- Improvement 23(4): 776-761. (In Farsi). <https://doi.org/10.22059/jci.2021.306872.2424>
- Khodabin G., Tahmasebi-Sarvestani Z., Shirani Rad A., Bakhshandeh E., Khayat Moghadam M., Kazemi S., Rafati Alashti M., Heidarzadeh A. 2022. Evaluation of the effects of zinc and manganese sulfate on the improvement of physiological characteristics and yield of rapeseed genotypes under late-season drought stress. *Journal of Crops Improvement* 24(4): 1217-1232. <http://doi.org/10.22059/jci.2022.334740.2645>
- Khosrowshahi Z.T., Ghassemi-Golezani K., Salehi-Lisar S.Y., Motafakkerzad R. 2020. Changes in antioxidants and leaf pigments of safflower (*Carthamus tinctorius* L.) affected by exogenous spermine under water deficit. *Biologia Futura* 71(3): 313-321. <https://doi.org/10.1007/s42977-020-00039-z>
- Kishor P.K., Sangam S., Amrutha R.N., Laxmi P.S., Naidu K.R., Rao K.S., Rao S., Reddy K.J., Theriappan P., Sreenivasulu N. 2005. Regulation of proline biosynthesis, degradation, uptake and transport in higher plants: its implications in plant growth and abiotic stress tolerance. *Current Science* 88(3): 424-438. <https://www.jstor.org/stable/24110209>
- Krüger H.G. 2002. Separately and simultaneously induced dark chilling and drought stress effects on photosynthesis, proline accumulation and antioxidant metabolism in soybean. *Journal of Plant Physiology* 159(10): 1077-1086. <https://doi.org/10.1078/0176-1617-00745>
- Kumar R.R., Karajol K., Naik G.R., 2011. Effect of polyethylene glycol induced water stress on physiological and biochemical responses in pigeonpea (*Cajanus cajan* L. Millsp.). *Recent Research in Science and Technology* 3(1): 148-152. <https://updatepublishing.com/journal/index.php/rrst/article/view/565>
- Li C.Z., Xie H.Y., Jin H.X. 2023. Extraction technology for manganese sulfate solution purification. *Engineering Chemistry* 1: 45-56. <https://doi.org/10.4028/p-6oe21m>
- Lidon F.C., Barreiro M.G., Ramalho J.C. 2004. Manganese accumulation in rice: implications for photosynthetic functioning. *Journal of Plant Physiology* 161(11): 1235-1244. <https://doi.org/10.1016/j.jplph.2004.02.003>
- MacAdam J.W., Nelson C.J., Sharp R.E. 1992. Peroxidase activity in the leaf elongation zone of tall fescue: I. Spatial distribution of ionically bound peroxidase activity in genotypes differing in length of the elongation zone. *Plant Physiology* 99(3): 872-878. <https://doi.org/10.1104/pp.99.3.872>
- Millaleo R., Reyes-Díaz M., Ivanov A.G., Mora M.L., Alberdi M. 2010. Manganese as essential and toxic element for plants: transport, accumulation and resistance mechanisms. *Journal of Soil Science and Plant Nutrition* 10(4): 470-481. <https://doi.org/10.4067/S0718-95162010000200008>
- Mishra S.S., Panda D. 2017. Leaf traits and antioxidant defense for drought tolerance during early growth stage in some popular traditional rice landraces from Koraput, India. *Rice Science* 24(4): 207-217. <https://doi.org/10.1016/j.rsci.2017.04.001>
- Moradi A., Piri R. 2018. Plant growth promoting rhizobacteria enhance salinity stress tolerance in Cumin (*Cuminum cyminum* L.) during germination stage. *Journal of Plant Process and Function* 6(22): 47-54. <https://dor.isc.ac/dor/20.1001.1.23222727.1396.6.22.8.2>
- Moradshahi A., Salehi Eskandari B., Kholdebarin B. 2004. Some physiological responses of canola (*Brassica napus* L.) to water deficit stress under laboratory conditions. *Iranian Journal of Science & Technology, Transaction* 28(1): 43-50. <https://doi.org/10.22099/ijsts.2004.2832>
- Movahhedy-Dehnavy M., Modarres-Sanavy S.A., Mokhtassi-Bidgoli A. 2009. Foliar application of zinc and manganese improves seed yield and quality of safflower (*Carthamus tinctorius* L.) grown under water deficit stress. *Industrial Crops and Products* 30(1): 82-92. <https://doi.org/10.1016/j.indcrop.2009.02.004>
- Muhammad I., Kolla M., Volker R., Günter N. 2015. Impact of nutrient seed priming on germination, seedling development, nutritional status and grain yield of maize. *Journal of Plant Nutrition* 38(12): 1803-1821. <https://doi.org/10.1080/01904167.2014.990094>
- Nakano Y., Asada K. 1987. Purification of ascorbate peroxidase in spinach chloroplast: in inactivation in ascorbate-depleted medium and reactivation by monodehydroascorbate radical. *Plant and Cell Physiology* 28(1): 131-140. <https://doi.org/10.1093/oxfordjournals.pcp.a077268>
- Piri R., Moradi A., Balouchi H., Salehi A. 2019. Improvement of cumin (*Cuminum cyminum*) seed performance under drought stress by seed coating and biopriming. *Scientia Horticulturae* 257: 108667. <https://doi.org/10.1016/j.scienta.2019.108667>
- Rasheed M.U., Malik A. 2022. Mechanism of drought stress tolerance in wheat. *Bulletin of Biological and Allied Sciences Research* 2022(1): 23. <https://doi.org/10.54112/bbasr.v2022i1.23>
- Sawhney V., Singh D.P. 2002. Effect of chemical desiccation at the post-anthesis stage on some physiological and biochemical changes in the flag leaf of contrasting wheat genotypes. *Field Crops Research* 77(1): 1-6. [https://doi.org/10.1016/S0378-4290\(01\)00192-7](https://doi.org/10.1016/S0378-4290(01)00192-7)
- Seifikalhor M., Niknam V., Aliniaiefard S., Didaran F., Tsaniklidis G., Fanourakis D., Teymoorzadeh M., Mousavi S.H., Bosacchi M., Li T. 2022. The regulatory role of γ -aminobutyric acid in chickpea plants depends on drought tolerance and water scarcity level. *Scientific Reports* 12(1): 7034. <https://doi.org/10.1038/s41598-022-10571-8>
- Setter T.L., Flannigan B.A., Melkonian J. 2001. Loss of kernel set due to water deficit and shade in maize: carbohydrate supplies, abscisic acid, and cytokinins. *Crop Science* 41(5): 1530-1540. <https://doi.org/10.2135/cropsci2001.4151530x>
- Seyed Ebrahimi F.S., Hassani Kumleh H., Alami A., Rezadoost M.H. 2018. Effect of drought stress on antioxidant activity in two sensitive and tolerant rapeseed (*Brassica napus*) cultivars in cell suspension culture. *Plant Production Technology* 9(2): 181-191. (In Farsi). <https://doi.org/10.22084/ppt.2017.2244>
- Shafiq S., Akram N.A., Ashraf M., García-Caparrós P., Ali O.M., Latef A.A. 2021. Influence of glycine betaine (natural and synthetic) on growth, metabolism and yield production of drought-stressed maize (*Zea mays* L.) plants. *Plants* 10(11): 2540. <https://doi.org/10.3390/plants10112540>
- Shoormij F., Mirlohi A., Saeidi G., Shirvani M. 2022. Combined foliar application of Zn and Fe increases grain micronutrient concentrations and alleviates water stress across diverse wheat

- species and ploidal levels. *Scientific Reports* 12(1): 20328. <https://doi.org/10.1038/s41598-022-24868-1>
- Siddiqui M.H., Al-Khaishany M.Y., Al-Qutami M.A., Al-Wahaibi M.H., Grover A., Ali H.M., Al-Wahaibi M.S., Bukhari N.A. 2015. Response of different genotypes of faba bean plant to drought stress. *International Journal of Molecular Sciences* 16(5): 10214-10227. <https://doi.org/10.3390/ijms160510214>
- Sobhanian H., Pahlavan S., Meyfour A. 2020. How does proteomics target plant environmental stresses in a semi-arid area? *Molecular Biology Reports* 47(4): 3181-3194. <https://doi.org/10.1007/s11033-020-05406-6>
- Soliman A.A., Mousa M.I., Ibrahim M.A., Baiomy K.A., Shaaban S.A., Shabana M.M., Mohamed E.N., Rehan M., Yu H., He Y. 2025. Agronomic and anatomic performance of some soybean genotypes under optimal and water-deficit conditions. *Frontiers in Plant Science* 16: 1575180. <https://doi.org/10.3389/fpls.2025.1575180>
- Su J., Zhang Y., Nie Y., Cheng D., Wang R., Hu H., Shen W. 2018. Hydrogen-induced osmotic tolerance is associated with nitric oxide-mediated proline accumulation and reestablishment of redox balance in alfalfa seedlings. *Environmental and Experimental Botany* 147: 249-260. <https://doi.org/10.1016/j.envexpbot.2017.12.022>
- Szabados L., Savaouré A. 2010. Proline: a multifunctional amino acid. *Trends in Plant Science* 15(2): 89-97. <https://doi.org/10.1016/j.tplants.2009.11.009>
- Theocharis A., Bordiec S., Fernandez O., Paquis S., Dhondt-Cordelier S., Baillieul F., Clément C., Barka E.A. 2012. Burkholderia phytofirmans PsJN primes *Vitis vinifera* L. and confers a better tolerance to low nonfreezing temperatures. *Molecular Plant-Microbe Interactions* 25(2): 241-249. <https://doi.org/10.1094/MPMI-05-11-0124>
- Wang X., Liu H., Yu F., Hu B., Jia Y., Sha H., Zhao H. 2019. Differential activity of the antioxidant defence system and alterations in the accumulation of osmolyte and reactive oxygen species under drought stress and recovery in rice (*Oryza sativa* L.) tillering. *Scientific Reports* 9(1): 8543. <https://doi.org/10.1038/s41598-019-44958-x>
- Waraich E.A., Ahmad R., Halim A., Aziz T. 2012. Alleviation of temperature stress by nutrient management in crop plants: a review. *Journal of Soil Science and Plant Nutrition* 12(2): 221-244. <https://doi.org/10.4067/S0718-95162012000200003>
- Zafari M., Ebadi A., Jahanbakhsh S., Sedghi M. 2020. Safflower (*Carthamus tinctorius*) biochemical properties, yield, and oil content affected by 24-epibrassinosteroid and genotype under drought stress. *Journal of Agricultural and Food Chemistry* 68(22): 6040-6047. <https://doi.org/10.1021/acs.jafc.9b06860>
- Zi X., Wang W., Zhou S., Zhou F., Rao D., Shen P., Fang S., Wu B. 2022. Prolonged drought regulates the silage quality of maize (*Zea mays* L.): alterations in fermentation microecology. *Frontiers in Plant Science* 13: 1075407. <https://doi.org/10.3389/fpls.2022.1075407>

HOW TO CITE THIS ARTICLE

Najafi G., Zarea M.J., Fazeli A., Zarei B. 202x. Drought Stress Amelioration in Sesame Cultivars: Unraveling the Combined Potential of MnSO₄ Seed Treatment and Foliar Nutrition. *Agrotechniques in Industrial Crops* x(x): xx-xx. [10.22126/ATIC.2026.12043.1212](https://doi.org/10.22126/ATIC.2026.12043.1212)