

Gene Expression Analysis of *IND*, *ALC*, *RPL*, and *FUL* in Reducing Seed Shattering in Flixweed (*Descurainia sophia*)

Yasoub Shiri^{*1} , Zeynab Mohkami¹ , Isa Khammari² 

¹Department of Agriculture and Plant Breeding, Agriculture Institute, Research Institute of Zabol, Zabol, Iran

²Department of Agronomy, Faculty of Agriculture, University of Zabol, Zabol, Iran

ARTICLE INFO

Original paper

Article history:

Received: xx xxx 202x

Revised: xx xxx 202x

Accepted: xx xxx 202x

Keywords:

Co-expression network

Harvest index

Indeterminate flowering

Plant hormone

ABSTRACT

The indeterminate growth habit and asynchronous pod maturation in flixweed (*Descurainia sophia*) lead to seed shattering and a consequent reduction in economic yield. Genes involved in the formation and differentiation of the dehiscence zone play a pivotal role by promoting or suppressing valve margin tissue identity. In this study, the expression of four key genes—*INDEHISCENT* (*IND*), *ALCATRAZ* (*ALC*), *REPLUMLESS* (*RPL*), and *FRUITFULL* (*FUL*)—was investigated under two plant hormone treatments (indole-3-butyric acid (IBA) and gibberellic acid (GA)), each applied at four different concentrations. Among these genes, *IND* and *ALC* promote seed shattering, whereas *RPL* and *FUL* function as negative regulators of pod dehiscence. In addition, six quantitative traits and their correlations with the expression of target genes were evaluated. The results showed that both IBA and GA influenced seed shattering by directly modulating the expression of genes responsible for the differentiation of the lignification layer (LL) and separation layer (SL). IBA at 40 mg L⁻¹ was identified as the most effective treatment for reducing seed shattering in flixweed. Correlation analyses revealed significant relationships ($r=0.31-0.58$, $p<0.05$) between the expression of *FUL* and *RPL* with seed yield and harvest index, whereas *IND* showed mainly negative correlations with thousand-seed weight ($r=-0.34$) and plant height ($r=-0.15$). Co-expression network analysis highlighted *HEC3*, a transcription factor involved in valve margin development, and *RPL* as central regulators in the genetic control of seed shattering. Plants treated with IBA 40 mg L⁻¹ exhibited the highest harvest index, increased expression of the seed-retention genes *FUL* and *RPL*, and reduced expression of the shattering-promoting genes *IND* and *ALC*. Although GA at 15 mg L⁻¹ represented the most effective treatment, IBA at 40 mg L⁻¹ consistently produced superior improvements in growth, yield, and seed shattering reduction.

DOI: [10.22126/ATIC.2026.12895.1240](https://doi.org/10.22126/ATIC.2026.12895.1240)

© The Author(s) 202x. Published by Razi University



1. Introduction

Flixweed (*Descurainia sophia*), also known by its synonyms *Sisymbrium sophia* and *Sisymbrium irio*, is an annual or biennial herb belonging to the family *Brassicaceae*. The plant can reach up to one meter in height, and its small, slightly elongated seeds are enclosed within slender pods that occur in two distinct colors (Lehnhoff *et al.*, 2025). A major agronomic challenge in flixweed cultivation is the asynchronous maturation of pods along the main stem, which leads to seed shattering and consequently lowers economic yield. In many plant species, the inflorescence meristem remains active after the transition from the

vegetative to the reproductive phase, continuously producing flowers and fruits. This growth pattern, known as indeterminate flowering/inflorescence, is characterized by a prolonged activity of the apical meristem, which keeps generating floral organs even after fruit set. In contrast, determinate inflorescence terminates after a fixed number of flowers, as the apical meristem differentiates into a terminal flower. Flixweed follows an indeterminate flowering pattern, typically blooming from late May to early July (Balanà *et al.*, 2019; Lehnhoff *et al.*, 2025). Genes involved in the formation and regulation of the pod dehiscence zone are critical for pod development and

* Corresponding author.

E-mail address: Yasub.shiri@uoaz.ac.ir

seed retention. Among these, *FUL*, *SHP1*, *SHP2*, *ALC*, *IND*, and *RPL* are of particular importance. The genes *SHATTERPROOF1* (*SHP1*), *SHATTERPROOF2* (*SHP2*), *IND*, and *ALC* enhance pod opening by promoting differentiation of the dehiscence zone, while *FUL* and *RPL* act antagonistically by suppressing shattering through their effects on valve margin cells. Thus, changes in the expression of these genes directly influence the degree of seed shattering in flaxweed. Reducing seed shattering must be achieved without compromising seed size or number, as pod retention alone does not guarantee maximum yield.

Pod dehiscence is regulated by a well-characterized genetic network in which several key transcription factors act in a coordinated manner. *FUL* functions as an upstream regulator that suppresses the expression of *SHP1* and *SHP2* in the valve tissues, thereby restricting dehiscence zone identity to the valve margin (Bemer et al., 2017; Hou et al., 2013). Whereas *SHP* genes do not exert reciprocal regulation on *FUL*, establishing a unidirectional regulatory hierarchy. This hierarchy provides a suitable framework to investigate whether exogenous application of IBA and GA can modulate pod dehiscence by altering the expression balance between upstream *FUL* and downstream *SHP*-*IND*/*ALC* components of the network. In full mutants, the natural dehiscence zone fails to differentiate. Lignification pattern analysis of the fruit margin revealed that, whereas wild-type siliques possess a single lignified valve cell layer, ful mutants show lignification throughout the entire valve tissue (Ferrándiz and Fourquin, 2014; Liu et al., 2016). In *Arabidopsis*, *SHP1* and *SHP2* promote the differentiation of valve margin cells by activating downstream regulators such as *IND* and *ALC*, which are essential for proper formation of the separation and lignification layers (Ehlers et al., 2016). Single mutations in *shp1* or *shp2* have little phenotypic effect, but double mutants produce non-dehiscent pods, indicating additive gene function. Unlike wild-type fruits, where lignification is observed in valve margin cells adjacent to the dehiscence zone, *shp1/shp2* mutants show reduced lignification. *FUL* negatively regulates *SHP1* and *SHP2* expression, whereas *SHP* genes have no reciprocal effect on *FUL*. These observations suggest that *SHP* genes are activated by *AGAMOUS* (*AG*) while *FUL* acts as a repressor of *SHP* expression (Deng et al., 2011; Dreni and Kater, 2014;

Hanano and Goto, 2011). *RPL* acts antagonistically to *SHP* genes by maintaining replum identity and spatially confining the dehiscence zone (Sun et al., 2023). Unlike *shp1/shp2* mutants that disrupt differentiation of the entire dehiscence zone, *alc* mutants exhibit normal external morphology and lignification patterns, affecting only a subset of valve margin tissues (Rajani and Sundaresan, 2001). Together, the *FUL*-*SHP*-*IND*/*ALC* regulatory module establishes precise spatial control over valve margin development, ultimately determining pod shattering behavior (Fig. 1).




Figure 1. Schematic representation of the genetic regulatory network controlling pod dehiscence in *Sisymbrium* spp. The transcription factors *SHP1/2* act as central regulators by activating *IND* and *ALC*, which in turn promote pod dehiscence through proper differentiation of the valve margin tissues. *FUL* negatively regulates *SHP1/2*, thereby restricting premature dehiscence, while *RPL* contributes to replum identity and limits the spatial expansion of the dehiscence zone. Arrows indicate activation or promotion, and blunt-ended lines indicate repression or restriction.

Given the importance of these genetic pathways, changes in the expression of pod-dehiscence genes strongly influence seed shattering (Zhang et al., 2020). However, emerging evidence suggests species-specific variations in gene function, indicating that functional validation in flaxweed is still required (Lotz et al., 2024). Plant hormones such as auxins are known to influence the expression of these bHLH genes and their downstream targets (Song et al., 2022). The *INDEHISCENT* (*IND*) gene encodes a basic helix–loop–helix (bHLH) transcription factor that specifies valve margin identity and plays a critical role in the differentiation of the lignification layer (LL) and separation layer (SL) (Liljegren et al., 2004). Conversely, *REPLUMLESS* (*RPL*) encodes a

homeodomain protein that functions similarly to *FUL* by negatively regulating valve margin identity genes within the replum (Xu et al., 2022). Auxins, through their roles in plant growth and tissue lignification, may affect the expression of these key regulators. Indole-3-butyric acid (IBA) was selected as the auxin treatment due to its greater chemical stability and more consistent physiological effects under exogenous application compared with IAA (Yang et al., 2025). Together, the combined evaluation of gibberellin and IBA provides an appropriate hormonal framework to investigate the regulation of seed shattering and yield-related traits in flaxweed. This study aimed to identify optimal concentrations of IBA and GA to enhance flaxweed yield through physiological regulation and modulation of key pod-dehiscence genes.

2. Materials and methods

This experiment was conducted in the research fields of the Research Institute of Zabol (UTM-x=377140.224, y=3412299.207) using a completely randomized design with three replications. Two plant growth regulators were applied: IBA at concentrations

of 10, 20, 30, and 40 mg L⁻¹, and GA at concentrations of 15, 30, 45, and 60 mg L⁻¹. These concentration ranges were chosen to encompass suboptimal to optimal levels reported in the literature, allowing the identification of dose-dependent responses in growth, yield, and seed shattering (Kalkali et al., 2023; Montazerinezhad et al., 2024; Paam et al., 2024; Ranjbar et al., 2024; Shiri et al., 2020). IBA and GA were applied in separate experimental treatments, allowing evaluation of their individual effects. Foliar sprays were performed at two growth stages—seedling and flowering—using the specified concentrations. Control plants were sprayed with distilled water. At physiological maturity, 20 plants were randomly selected from each plot to measure quantitative traits.

To examine the expression of the four key genes (*IND*, *ALC*, *RPL*, and *FUL*)—among which *IND* and *ALC* promote seed shattering, while *RPL* and *FUL* suppress it—four pairs of gene-specific primers were designed based on mRNA sequences retrieved from the NCBI database, using Primer3 Plus and CLC Main Workbench v7 (Table 1). *Actin7* was used as the reference gene.

Table 1. Sequences of primers used in the PCR reaction. The ACT7 gene was used as a reference gene.

Gene name	Gene ID	Primers 5'-3'		Product length (bp)
FUL	XM_024151164.1	Fw	TGGTCATTTTCAGGGTTGTCA	155
		Rv	ACCAGACCTTCGCTTTGAGA	
RPL	XM_006398603.2	Fw	CGGACCAGCTTCAAGTTAGC	134
		Rv	TTGACCAGCAGAACAACAAGC	
IND	XM_006396147.2	Fw	AGTTCTTGAAACGGCAGGTG	188
		Rv	CAATCGTCGATCGAAATCCT	
ALC	XM_006393847.2	Fw	TGAGGCCTTACAGCCAGAGT	157
		Rv	GCCGAATCATCACATCACTG	
Actin7	XM_006399401.2	Fw	TTCCCAAGTATTGTTGGTCGTC	168
		Rv	CCAGATCTTCTCCATGTCATCC	

Total RNA was extracted following the method of Japelaghi (Japelaghi et al., 2011). RNA quality and quantity were assessed by spectrophotometry and horizontal gel electrophoresis, and suitable samples were used for cDNA synthesis. Using a thermal cycler with gradient temperature settings, the optimal annealing temperature for each gene was determined. Quantitative real-time PCR was subsequently performed (in three biological and three technical replicates) with a Rotor-Gene 3000 system using a 2× Taq Master Mix PCR kit (CinaGene, Iran). Relative expression levels were calculated using the $\Delta\Delta CT$ method (Livak and Schmittgen, 2001). To construct the

gene co-expression network, interaction data were retrieved from the STRING v11 database (Szklarczyk et al., 2023). Co-expression relationships were inferred using the co-expression evidence channel implemented in STRING v11. Network visualization was carried out using Cytoscape v3.7.1 software (Ono et al., 2025).

Quantitative traits, including harvest index, number of dehisced pods, seed yield, thousand-seed weight, plant height, and number of seeds per pod, were analyzed. The experiment was conducted as a completely randomized design (CRD) with three biological replicates. Data were subjected to analysis of variance (ANOVA) using SAS software (v. 9), and

mean comparisons were performed using an appropriate post-hoc test at $p < 0.05$. Correlations between gene expression levels and quantitative traits were evaluated using Pearson's correlation coefficient.

An additional parameter measured was the proportion of dehiscent pods per plant. Because the total pod number varied among plants, the proportional measure of dehiscent pods was calculated using the ratio of dehiscent pods to total pods (Equation 1).

$$(1) \quad x = \frac{ab}{c}$$

Where x represents the ratio of dehiscent pods to the total number of pods, a is the total number of examined plants (20 plants per treatment), b is the total number of split pods, and c is the total number of examined pods.

3. Results and discussion

While new pods continue to form along the upper parts of the flowering stem, the lower pods mature and eventually dehisce, releasing their seeds (Fig. 2). Such asynchronous maturation leads to substantial seed loss—up to 30% of potential yield (Takahashi et al., 2020)—causing not only reduced economic return but also the emergence of volunteer seedlings in subsequent seasons and complicating harvest timing.




Figure 2. Over a period of 3 to 4 weeks, Flixweed continuously produces new pods while the older pods split open and release their seeds. A cross-section of two dehiscent pods and one intact pod clearly shows the differentiated cellular layers in the pod wall. Four key genes, by regulating the differentiation of these cells, control the process of pod dehiscence.

Plant hormones can modify pod dehiscence by influencing the expression of genes associated with pod

opening. They also affect other quantitative traits such as plant height, seed number per pod, and thousand-seed weight. Therefore, reducing seed shattering must be achieved without compromising seed size or number, as pod retention alone does not guarantee maximum yield.

3.1. Quantitative trait analysis

Six quantitative traits—plant height, seed number per pod, thousand-seed weight, seed yield, and harvest index—were analyzed to identify the most effective hormone and concentration (Fig. 3). IBA treatments ($10\text{--}40 \text{ mg L}^{-1}$) significantly increased plant height compared with the control, with the greatest effect observed at 40 mg L^{-1} (Fig. 3a). By contrast, GA treatments, particularly at 60 mg L^{-1} , caused a marked reduction in plant height. These findings are consistent with previous reports describing auxins as strong stimulators of shoot elongation and vegetative growth (Lestari et al., 2024).

Gibberellins are generally recognized as key regulators of internode elongation (Dai et al., 2025); however, plant responses to GA are non-linear, and high concentrations can trigger negative feedback mechanisms such as induction of GA-catabolic genes (e.g., GA2ox) or hormonal imbalance with auxins, resulting in growth inhibition (Hedden and Sponsel, 2015). The inhibitory effect of high GA concentrations was also evident in seed number per pod (Fig. 3b). Similar non-linear responses have been reported in other species: for example, in okra, GA application increased plant height but had little effect on flower initiation or pod formation (Mohammadi et al., 2014); in soybean, pod set is regulated by GA biosynthesis during post-flowering vegetative growth (Taniguchi et al., 2018); and in cauliflower (*Brassica oleracea*), GA at 200 ppm slightly increased seed number, while higher concentrations significantly reduced it (Prodhan et al., 2022).

Collectively, these studies confirm the dose-dependent and potentially inhibitory effects of GA (Abdel-Mola et al., 2022; Kalkali et al., 2023; Montazerinezhad et al., 2024; Paam et al., 2024; Ranjbar et al., 2024; Shiri et al., 2018, Shiri et al., 2020). It should be noted that plant responses to gibberellic acid are highly species-specific and may vary depending on developmental and physiological context.




Figure 3. Graphs of the quantitative traits examined under IBA and GA treatments in flaxweed. The traits are: a. plant height, b. number of seeds per pod (Average number counted), c. thousand-seed weight, d. grain yield, e. harvest index, and f. number of open pods per plant (Average number counted). The IBA treatment had a greater enhancing effect on growth and yield-related indices compared to GA.

Although most previous studies have focused on the role of IBA in rooting and plant establishment, some evidence supports its positive influence on seed yield. For example, IBA spraying improved seed production in onion by increasing seed number per fruit (Gabri *et al.*, 2024), which agrees with the present results. Overall, the growth and yield indices clearly demonstrate that IBA treatments outperformed GA treatments (Fig. 3). Similar positive effects of IBA on vegetative growth and yield have been reported in rapeseed, where IBA enhanced stem length, assimilate accumulation, and final seed yield (Li *et al.*, 2024; Li *et al.*, 2018). In contrast, GA responses are less predictable and depend strongly on concentration,

timing, and application method; in some cases, GA sprays negatively affect seed production by disturbing floral organ development (Castro-Camba *et al.*, 2022).

Hormone treatments significantly reduced pod dehiscence compared with the control, with the greatest reduction observed in plants treated with IBA 40 mg L⁻¹. This finding indicates that optimal IBA concentration effectively prevents pod splitting—a key factor in minimizing yield loss. Reducing seed shattering must be achieved without compromising seed size or number, as pod retention alone does not guarantee maximum yield. Auxins and their derivatives are known to increase adhesion of valve margin cells and interfere with tissue differentiation and lignification

processes, thereby enhancing pod integrity and seed retention (Dong and Wang, 2015; Liu et al., 2024). Auxin treatments likely reduce pod dehiscence by reinforcing valve margin cell adhesion through coordinated transcriptional regulation. The upregulation of *FUL* and *RPL* suppresses the expression of the bHLH genes *IND* and *ALC*, which are required for differentiation and separation of the valve margin. This regulatory shift limits the formation of a functional separation layer, thereby reducing cell wall loosening and preventing premature pod opening.

3.2. Expression analysis of target genes

The fruits of *Arabidopsis*, rapeseed, and other Brassicaceae members are siliques composed of two valves attached to a central replum through a specialized valve margin tissue. The valve margin contains two distinct layers: a lignification layer (LL), which provides mechanical tension, and a separation layer (SL), which secretes polygalacturonases to degrade cell walls and allow cell separation (Arnaud et al., 2010). Genes *IND* and *ALC* promote differentiation of LL and SL, respectively, and their increased expression facilitates pod opening. Conversely, *FUL* and *RPL* suppress the action of *IND* and *ALC*, thereby reducing pod dehiscence (Singh et al., 2024).

Previous studies have shown that auxin and gibberellin independently influence the expression of genes involved in LL and SL differentiation. In *Arabidopsis*, *IND* acts as a central coordinator of multiple hormonal pathways (Simonini et al., 2017). In the present study, we observed that *IND* expression responded to exogenous IBA and GA treatments, although not in a strictly dose-dependent manner. This

responsiveness suggests that *IND* may act as a downstream convergence point for hormone-mediated regulatory signals in flaxweed. While our data do not directly dissect the molecular crosstalk mechanisms, the observed transcriptional modulation of *IND* under hormone treatments supports its potential role as an integrative node within the regulatory network. GA biosynthesis, enhanced by *IND*, is required for silique opening and seed dispersal (van Gelderen et al., 2016). Localized auxin production is also essential for SL differentiation (Li et al., 2019). Exogenous hormone treatments can therefore supply sufficient hormone levels to downregulate the expression of genes required for endogenous hormone biosynthesis, but plant responses remain highly non-linear (Maity et al., 2021).

As shown in Fig. 4, IBA treatments increased the expression of *FUL* and *RPL* in a concentration-dependent manner, consistent with the observed reduction in pod dehiscence. In contrast to *FUL* and *RPL*, whose expression showed a clear dose-dependent increase under rising IBA concentrations and was consistently associated with a reduction in pod shattering, the expression patterns of *IND* and *ALC* did not follow this monotonic trend. Instead, their transcriptional responses varied across hormone concentrations and were not directly aligned with the observed decrease in the number of dehiscent pods. Moya-Cuevas et al. (2024), through transcriptomic modeling analysis, confirmed the repressive roles of *FUL* and *RPL* on the *SHP/IND/ALC* module and the promotive roles of *IND* and *ALC* in SL formation, supporting the present findings (Moya-Cuevas et al., 2024).




Figure 4. Comparison of the mean relative expression of genes involved in the pod development of Flaxweed under the PGRs treatments of GA and IBA. By upregulation of *FUL/RPL* and downregulation of *IND/ALC*, IBA at 40 mg L⁻¹ emerged as the most effective treatment for reducing seed shattering in flaxweed.

3.3. Correlations between gene expression and quantitative traits

Correlation analysis revealed that *FUL* and *RPL* showed the strongest positive associations with seed yield and harvest index, while *IND* exhibited negative correlations with thousand-seed weight and plant height (Fig. 5). These relationships underscore the potential of *FUL* and *RPL* as enhancers of yield and harvesting efficiency, and the limiting role of *IND*. Similar patterns have been reported in Arabidopsis and other Brassicaceae species, where *FUL* promotes fruit tissue differentiation and seed filling (Di Marzo et al., 2020), and *RPL* enhances silique growth and seed production capacity (Roeder et al., 2003). Conversely, *IND* is a key activator of pod opening and seed dispersal, and *ALC* participates in seed release and fruit maturation (Ivarson et al., 2023). These results suggest that suppression of *IND* expression could be a promising breeding strategy to reduce seed shattering and improve yield stability in flaxweed.




Figure 5. Heatmap diagram showing the correlation of the expression of four genes, *FUL*, *RPL*, *IND* and *ALC* with changes in the quantitative traits in the flaxweed. The color intensity from blue (negative correlation) to red (positive correlation) indicates the direction and magnitude of the correlation coefficient. As can be seen, the expression of *RPL* and *FUL* has the highest positive correlation with harvest index and grain yield, while thousand-grain weight shows a significant negative correlation with most genes.

Interestingly, thousand-seed weight showed a negative correlation with *IND* and *ALC* expression, which are known promoters of pod shattering. Although this relationship appears counterintuitive, it may reflect a developmental trade-off. Elevated expression of *IND* and *ALC* could accelerate valve margin differentiation and separation layer formation, potentially reallocating assimilates toward structural remodeling processes rather than seed filling. Alternatively, higher activity of these shattering-related genes may be associated with earlier pod maturation,

thereby shortening the effective seed-filling period. Further functional analyses are required to clarify this relationship.

3.4. Gene co-expression network analysis

Network reconstruction revealed a central role for *HEC3* in flaxweed pod maturation (Fig. 6). Topological analysis using betweenness centrality (BC) and closeness centrality (CC) identified *HEC3* and *RPL* as hub genes with the highest BC value (0.0714) (Table 2).




Figure 6. Gene network of the studied genes involved in seed shattering in flaxweed. The size of each node represents the CC value (larger nodes indicate higher values), and the color of each node represents the BC value (nodes with higher values are closer to red, while lower values are closer to blue). Similarly, thicker and more reddish edges indicate higher Edge Betweenness (EB) values. The network topology suggests that the *HEC3* gene plays an important role in the maturation process of flaxweed siliques.

Table 2. Topological analysis of coexpression network of genes involved in Flaxweed shattering seed

Name	Edge count	Betweenness centrality	Closeness centrality
HEC3	4	0.071429	1
RPL	4	0.071429	0.666667
ACI1	3	0	1
SPT	3	0	0
FUL	2	0	0.75
ALC	1	0	0
SHP1	1	0	0.5
SHP2	1	0	0.5
IND	1	0	1

The prominent network position of *HEC3* is noteworthy. Although *ACI1* and *IND* also showed high CC values, *HEC3* and *RPL* were distinguished by their simultaneous prominence across all evaluated parameters. Therefore, our interpretation emphasizes these two genes as principal hubs within the network. Members of the *HECATE* (*HEC1/2/3*) family are bHLH transcription factors expressed in the female reproductive tissues of the gynoecium (Arias-Pérez et al., 2025). These regulatory roles have been associated

with the establishment of transmission tissues and hormone gradients required for activation of *IND* and *ALC* at the valve margin (Schuster et al., 2015). In the present study, this connection is interpreted in light of previous findings, as hormone gradients were not directly quantified. *HEC3* was not included as an initial candidate gene and, to our knowledge, has not been directly implicated in pod shattering in previous studies. However, it emerged as a central hub in the unbiased co-expression network constructed from our dataset. Given its prominent network connectivity, *HEC3* may represent a previously unrecognized regulatory component associated with pod dehiscence in flaxweed. This observation should be considered hypothesis-generating and requires further functional validation. Although the genetic network controlling pod dehiscence has been extensively characterized in Arabidopsis, its functional relevance in flaxweed remains largely unexplored. Given the distinct growth habit and agronomic importance of flaxweed, functional validation of this network in this species is required.

4. Conclusion

Hormonal regulation significantly influences pod dehiscence and, consequently, the final yield of flaxweed (*Descurainia sophia*). In this study, two plant growth regulators—indole-3-butyric acid (IBA) at concentrations of 10, 20, 30, and 40 mg L⁻¹ and gibberellic acid (GA) at concentrations of 15, 30, 45, and 60 mg L⁻¹—were applied to evaluate their effects on pod shattering and associated gene expression. Five key agronomic traits (plant height, seed number per pod, thousand-seed weight, seed yield, and harvest index) were analyzed to determine the most effective treatment. The results demonstrated that both IBA and GA independently modulated the expression of genes controlling the differentiation of the lignification layer (LL) and separation layer (SL), which are essential for pod opening. Overall, IBA treatments produced stronger positive effects on growth and yield indices compared with GA. Correlation analyses revealed that *FUL* and *RPL* had the highest positive associations with seed yield and harvest index, whereas *IND* exhibited negative correlations with thousand-seed weight and plant height. Considering the combined evidence from yield indices, quantitative trait analyses, pod dehiscence rates, and the expression patterns of

key genes, IBA at 40 mg L⁻¹ emerged as the most effective treatment for reducing seed shattering in flaxweed. Plants treated with IBA 40 mg L⁻¹ achieved the highest harvest index, showed elevated expression of the seed-retention genes *FUL* and *RPL*, and exhibited reduced expression of the shattering-promoting genes *IND* and *ALC*. Overall, IBA at 40 mg L⁻¹ proved to be the most effective treatment in enhancing yield-related traits and reducing silique shattering. These findings suggest that the application of IBA at this concentration represents a practical and viable strategy to mitigate yield loss in flaxweed cultivation. The strong edge between *HEC3* and *RPL*, along with their high CC values, suggests that these genes are suitable targets for expression manipulation or transient genetic interference (e.g., RNAi/CRISPRi) to further elucidate the regulatory mechanisms of seed shattering in flaxweed. Taken together, the network topology analysis, highlighting *HEC3* and *RPL* as central regulatory nodes, aligns with the observed transcriptional modulation and physiological improvements under IBA treatment, collectively linking molecular coordination with reduced pod shattering and enhanced yield performance in flaxweed.

Conflict of interests

All authors declare no conflict of interest.

Ethics approval and consent to participate

No humans or animals were used in the present research. The authors have adhered to ethical standards, including avoiding plagiarism, data fabrication, and double publication.

Consent for publications

All authors read and approved the final manuscript for publication.

Availability of data and material

All the data are embedded in the manuscript.

Authors' contributions

All authors had an equal role in study design, work, statistical analysis and manuscript writing.

Informed consent

The authors declare not to use any patients in this research.

Funding/Support

This research was supported by the Research Institute of Zabol, Project code: PR-RIOZ-1403-4183-1.

References

- Abdel-Mola M.A.M., Kenawy A.G.M., Ali S.K., Ayyat A.M. 2022. Influence of foliar application of gibberellic acid and liquorice root extract on growth, volatile oil yield productivity and antimicrobial activity in geranium (*Pelargonium graveolens* L. her.) plants. *Scientific Journal of Flowers and Ornamental Plants* 9(4): 249-271. <https://doi.org/10.21608/sjfor.2022.272767>
- Arias-Pérez E., Luna-García V., Bernal-Gallardo J.J., de Folter S. 2025. Identifying new players of gynoecium development using tissue-specific transcriptome data of *Arabidopsis*. *Planta* 262(3): 67. <https://doi.org/10.1007/s00425-025-04784-0>
- Arnaud N., Girin T., Sorefan K., Fuentes S., Wood T.A., Lawrenson T., Sablowski R., Østergaard L. 2010. Gibberellins control fruit patterning in *Arabidopsis thaliana*. *Genes & Development* 24(19): 2127-2132. <https://doi.org/10.1101/gad.593410>
- Balanà V., Martínez-Fernández I., Sato S., Yanofsky M.F., Ferrándiz C. 2019. Inflorescence meristem fate is dependent on seed development and FRUITFULL in *Arabidopsis thaliana*. *Frontiers in Plant Science* 10: 1622. <https://doi.org/10.3389/fpls.2019.01622>
- Bemer M., van Mourik H., Muiño J.M., Ferrándiz C., Kaufmann K., Angenent G.C. 2017. FRUITFULL controls SAUR10 expression and regulates *Arabidopsis* growth and architecture. *Journal of Experimental Botany* 68(13): 3391-3403. <https://doi.org/10.1093/jxb/erx184>
- Castro-Camba R., Sánchez C., Vidal N., Vielba J.M. 2022. Plant development and crop yield: the role of gibberellins. *Plants (Basel)* 11(19): 2650. <https://doi.org/10.3390/plants11192650>
- Dai L., Zhao X., Liu S., Keerthana K., Vijayakanth V., Zhi Y., Chen M., Que F., Ramakrishnan M., Ahmad Z., Wei Q. 2025. Gibberellin-mediated internode elongation in grasses with a focus on bamboo: molecular pathways and regulatory networks. *Frontiers in Plant Science* 16: 1665328. <https://doi.org/10.3389/fpls.2025.1665328>
- Deng W., Ying H., Helliwell C.A., Taylor J.M., Peacock W.J., Dennis E.S. 2011. FLOWERING LOCUS C (FLC) regulates development pathways throughout the life cycle of *Arabidopsis*. *Proceedings of the National Academy of Sciences* 108(16): 6680-6685. <https://doi.org/10.1073/pnas.1103175108>
- Di Marzo M., Herrera-Ubaldo H., Caporali E., Novák O., Strnad M., Balanà V., Ezquer I., Mendes M.A., de Folter S., Colombo L. 2020. SEEDSTICK controls *Arabidopsis* fruit size by regulating cytokinin levels and FRUITFULL. *Cell Reports* 30(8): 2846-2857.e3. <https://doi.org/10.1016/j.celrep.2020.01.101>
- Dong Y., Wang Y.Z. 2015. Seed shattering: from models to crops. *Frontiers in Plant Science* 6: 476. <https://doi.org/10.3389/fpls.2015.00476>
- Dreni L., Kater M.M. 2014. MADS reloaded: evolution of the AGAMOUS subfamily genes. *New Phytologist* 201(3): 717-732. <https://doi.org/10.1111/nph.12555>
- Ehlers K., Bhide A.S., Tekleyohans D.G., Wittkop B., Snowdon R.J., Becker A. 2016. The MADS box genes ABS, SHP1, and SHP2 are essential for the coordination of cell divisions in ovule and seed coat development and for endosperm formation in *Arabidopsis thaliana*. *PLOS ONE* 11(10): e0165075. <https://doi.org/10.1371/journal.pone.0165075>
- Ferrándiz C., Fourquin C. 2014. Role of the FUL-SHP network in the evolution of fruit morphology and function. *Journal of Experimental Botany* 65(16): 4505-4513. <https://doi.org/10.1093/jxb/ert479>
- Gabri C.G., Soto V.C., Ibañez A.A., Galmarini C.R. 2024. Foliar application of boron, calcium, and indole-3-butyric acid on onion plants to improve seed production. *Italian Journal of Agronomy* 19(2): 100008. <https://doi.org/10.1016/j.ijagro.2024.100008>
- Hanano S., Goto K. 2011. *Arabidopsis* TERMINAL FLOWER1 is involved in the regulation of flowering time and inflorescence development through transcriptional repression. *Plant Cell* 23(9): 3172-3184. <https://doi.org/10.1105/tpc.111.088641>
- Hedden P., Sponsel V. 2015. A century of gibberellin research. *Journal of Plant Growth Regulation* 34(4): 740-760. <https://doi.org/10.1007/s00344-015-9546-1>
- Hou K., Wu W., Gan S.S. 2013. SAUR36, a small auxin up RNA gene, is involved in the promotion of leaf senescence in *Arabidopsis*. *Plant Physiology* 161(2): 1002-1009. <https://doi.org/10.1104/pp.112.212787>
- Ivarson E., Ahlman A., Englund J.E., Lager I., Zhu L.H. 2023. Downregulation of the *INDEHISCENT* gene by RNAi resulted in desired pod shatter reduction of *Lepidium campestre* in subsequent generations. *International Journal of Molecular Sciences* 24(21): 15943. <https://doi.org/10.3390/ijms242115943>
- Japelaghi R.H., Haddad R., Garoosi G.A. 2011. Rapid and efficient isolation of high quality nucleic acids from plant tissues rich in polyphenols and polysaccharides. *Molecular Biotechnology* 49(2): 129-137. <https://doi.org/10.1007/s12033-011-9384-8>
- Kalkali N., Emamjomeh A., Shiri Y., Solouki M. 2023. Investigation of hormonal treatments in physiological alterations of Sistan Yaghooti grapes. *Applied Biology* 36(1): 159-180. (In Farsi). <https://doi.org/10.22051/jab.2022.40942.1497>
- Lehnhoff E.A., Schutte B.J., Rashid A., Sanogo S. 2025. *Descurainia sophia* (flixweed): a weed with many uses and ecological roles. In: *Agroecology of Edible Weeds and Noncrop Plants: Ecology and Socioeconomic Potential of the Associated Plant Biodiversity*. Elsevier, (pp. 209-224). <https://doi.org/10.1016/B978-0-443-16076-9.00013-5>
- Lestari D., Cahyaningrum D.T., Mayasari F., Suwarsono M.D. 2024. Application of indole butyric acid (IBA) to improve the quality of MD2 pineapple seedlings from crown shoots and sapling shoots. *IOP Conference Series: Earth and Environmental Science* 1338(1): 012012. <https://doi.org/10.1088/1755-1315/1338/1/012012>

- Li J.H., Feng N.J., Zheng D.F., Du X.L., Wu J.S., Wang X. 2024. Regulation of seed soaking with indole-3-butyric acid potassium salt (IBA-K) on rapeseed (*Brassica napus* L.) seedlings under NaCl stress. BMC Plant Biology 24(1): 904. <https://doi.org/10.1186/s12870-024-05586-4>
- Li S.W., Zeng X.Y., Leng Y., Feng L., Kang X.H. 2018. Indole-3-butyric acid mediates antioxidative defense systems to promote adventitious rooting in mung bean seedlings under cadmium and drought stresses. Ecotoxicology and Environmental Safety 161: 332-341. <https://doi.org/10.1016/j.ecoenv.2018.06.003>
- Li X.R., Vroomans R.M.A., Fox S., Grieneisen V.A., Østergaard L., Marée A.F.M. 2019. Systems biology approach pinpoints minimum requirements for auxin distribution during fruit opening. Molecular Plant 12(6): 863-878. <https://doi.org/10.1016/j.molp.2019.05.003>
- Liljegren S.J., Roeder A.H., Kempin S.A., Gremski K., Østergaard L., Guimil S., Reyes D.K., Yanofsky M.F. 2004. Control of fruit patterning in Arabidopsis by INDEHISCENT. Cell 116(6): 843-853. [https://doi.org/10.1016/S0092-8674\(04\)00217-X](https://doi.org/10.1016/S0092-8674(04)00217-X)
- Liu J., Wang W., Mei D., Wang H., Fu L., Liu D., Li Y., Hu Q. 2016. Characterizing variation of branch angle and genome-wide association mapping in rapeseed (*Brassica napus* L.). Frontiers in Plant Science 7: 21. <https://doi.org/10.3389/fpls.2016.00021>
- Liu L., Javed H.H., Hu Y., Luo Y.Q., Peng X., Wu Y.C. 2024. Research progress and mitigation strategies for pod shattering resistance in rapeseed. PeerJ 12: e18105. <https://doi.org/10.7717/peerj.18105>
- Livak K.J., Schmittgen T.D. 2001. Analysis of relative gene expression data using real-time quantitative PCR and the 2- $\Delta\Delta$ CT method. Methods 25(4): 402-408. <https://doi.org/10.1006/meth.2001.1262>
- Lotz D., Rössner L.H., Ehlers K., Kong D., Rössner C., Rupp O., Becker A. 2024. Conservation of the dehiscence zone gene regulatory network in dicots and the role of the SEEDSTICK ortholog of California poppy (*Eschscholzia californica*) in fruit development. EvoDevo 15(1): 16. <https://doi.org/10.1186/s13227-024-00236-0>
- Maity A., Lamichaney A., Joshi D.C., Bajwa A., Subramanian N., Walsh M., Bagavathiannan M. 2021. Seed shattering: a trait of evolutionary importance in plants. Frontiers in Plant Science 12: 657773. <https://doi.org/10.3389/fpls.2021.657773>
- Mohammadi G., Khah E.M., Petropoulos S.A., Chachalis D.B., Akbari F., Yarsi G. 2014. Effect of gibberellic acid and harvesting time on the seed quality of four okra cultivars. Journal of Agricultural Science 6(7): 200-211. <https://doi.org/10.5539/jas.v6n7p200>
- Montazerinezhad S., Solouki M., Emamjomeh A., Kavousi K., Taheri A., Shiri Y. 2024. Transcriptomic analysis of alternative splicing events for different stages of growth and development in Sistan Yaghooti grape clusters. Gene 896: 148030. <https://doi.org/10.1016/j.gene.2023.148030>
- Moya-Cuevas J., Ortiz-Gutiérrez E., López-Sánchez P., Simón-Moya M., Ballester P., Álvarez-Buylla E.R., Ferrándiz C. 2024. A model for the gene regulatory network along the Arabidopsis fruit medio-lateral axis: rewiring the pod shatter process. Plants 13(20): 2927. <https://doi.org/10.3390/plants13202927>
- Ono K., Fong D., Gao C., Churas C., Pillich R., Lenkiewicz J., Pratt D., Pico A.R., Hanspers K., Xin Y., Morris J., Kucera M., Franz M., Lopes C., Bader G., Ideker T., Chen J. 2025. Cytoscape Web: bringing network biology to the browser. Nucleic Acids Research 53(W1): W203-W212. <https://doi.org/10.1093/nar/gkaf365>
- Paam H., Emamjomeh A., Shiri Y. 2024. Effects of gibberellic acid and auxin on expression of genes involved in flaxweed flowering. Iranian Journal of Plant Physiology 14(3): 5087-5095. <https://doi.org/10.7155/ijpp.2024.1025881>
- Prodhan M.M., Sarker U., Hoque M.A., Biswas M.S., Ercisli S., Assouguem A., Ullah R., Almutairi M.H., Mohamed H.R.H., Najda A. 2022. Foliar application of GA3 stimulates seed production in cauliflower. Agronomy 12(6): 1394. <https://doi.org/10.3390/agronomy12061394>
- Rajani S., Sundaresan V. 2001. The Arabidopsis myc/bHLH gene ALCATRAZ enables cell separation in fruit dehiscence. Current Biology 11(24): 1914-1922. [https://doi.org/10.1016/S0960-9822\(01\)00593-0](https://doi.org/10.1016/S0960-9822(01)00593-0)
- Ranjbar S., Emamjomeh A., Ebrahimi M., Ghorbani A., Ebrahimie E., Shiri Y. 2024. Comparative analysis of bioinformatic tools to predict and quantify active circular RNAs during grape cluster development. Biologia Plantarum 68: 1-11. <https://doi.org/10.32615/bp.2023.038>
- Roeder A.H.K., Ferrándiz C., Yanofsky M.F. 2003. The role of the REPLUMLESS homeodomain protein in patterning the Arabidopsis fruit. Current Biology 13(18): 1630-1635. <https://doi.org/10.1016/j.cub.2003.08.027>
- Schuster C., Gaillochet C., Lohmann J.U. 2015. Arabidopsis HECATE genes function in phytohormone control during gynoecium development. Development 142(19): 3343-3350. <https://doi.org/10.1242/dev.120444>
- Shiri Y., Solouki M., Ebrahimie E., Emamjomeh A., Zahiri J. 2018. Unraveling the transcriptional complexity of compactness in Sistan grape cluster. Plant Science 270: 198-208. <https://doi.org/10.1016/j.plantsci.2018.02.011>
- Shiri Y., Solouki M., Ebrahimie E., Emamjomeh A., Zahiri J. 2020. Gibberellin causes wide transcriptional modifications in the early stage of grape cluster development. Genomics 112(1): 820-830. <https://doi.org/10.1016/j.ygeno.2019.05.022>
- Simonini S., Bencivenga S., Trick M., Østergaard L. 2017. Auxin-induced modulation of ETTIN activity orchestrates gene expression in Arabidopsis. The Plant Cell 29(8): 1864-1882. <https://doi.org/10.1105/tpc.17.00389>
- Singh A.K., Pandey S.K., Singh B.P., Chauhan R.K. 2024. Benefits of using plant growth regulators to enhance fruit production. International Journal of Advanced Biochemistry Research 8(9): 74-78. <https://doi.org/10.33545/26174693.2024.v8.i9b.2080>
- Song Y., Li S., Sui Y., Zheng H., Han G., Sun X., Yang W., Wang H., Zhuang K., Kong F., Meng Q., Sui N. 2022. SbbHLH85, a bHLH member, modulates resilience to salt stress by regulating root hair growth in sorghum. Theoretical and Applied Genetics 135(1): 201-216. <https://doi.org/10.1007/s00122-021-03960-6>
- Sun W., Chen Y., Zeng J., Li C., Yao M., Liu M., Ma Z., Huang L., Yan J., Zhan J., Chen H., Bu T., Tang Z., Li Q., Wu Q., Hou J., Huang Y. 2023. The Tartary buckwheat bHLH gene ALCATRAZ contributes to silique dehiscence in Arabidopsis

- thaliana*. Plant Science 333: 111733. <https://doi.org/10.1016/j.plantsci.2023.111733>
- Szklarczyk D., Kirsch R., Koutrouli M., Nastou K., Mehryary F., Hachilif R., Gable A.L., Fang T., Doncheva N.T., Pyysalo S., Bork P., Jensen L.J., von Mering C. 2023. The STRING database in 2023: protein-protein association networks and functional enrichment analyses for any sequenced genome of interest. Nucleic Acids Research 51(D1): D638-D646. <https://doi.org/10.1093/nar/gkac1000>
- Takahashi Y., Kongjaimun A., Muto C., Kobayashi Y., Kumagai M., Sakai H., Satou K., Teruya K., Shiroma A., Shimoji M., Hirano T., Isemura T., Saito H., Baba-Kasai A., Kaga A., Somta P., Tomooka N., Naito K. 2020. Same locus for non-shattering seed pod in two independently domesticated legumes, *Vigna angularis* and *Vigna unguiculata*. Frontiers in Genetics 11: 748. <https://doi.org/10.3389/fgene.2020.00748>
- Taniguchi T., Murayama N., Hasegawa M., Nakagawa A.C.S., Tanaka S., Zheng S.H., Hamaoka N., Iwaya-Inoue M., Ishibashi Y. 2018. Vegetative growth after flowering through gibberellin biosynthesis regulates pod setting rate in soybean (*Glycine max* (L.) Merr.). Plant Signaling & Behavior 13(8): e1473668. <https://doi.org/10.1080/15592324.2018.1473668>
- van Gelderen K., van Rongen M., Liu A., Otten A., Offringa R. 2016. An INDEHISCENT-controlled auxin response specifies the separation layer in early Arabidopsis fruit. Molecular Plant 9(6): 857-869. <https://doi.org/10.1016/j.molp.2016.03.005>
- Xu M., Wang X., Liu J., Jia A., Xu C., Deng X.W., He G. 2022. Natural variation in the transcription factor REPLUMLESS contributes to both disease resistance and plant growth in Arabidopsis. Plant Communications 3(5): 100351. <https://doi.org/10.1016/j.xplc.2022.100351>
- Yang B., Luo Z., Zhu X., Ji Y., Ma Q., Yan F. 2025. Research progress on the application of plant growth regulators in the rapid propagation of jujube by in vitro culture. Plants 14(19): 3012. <https://doi.org/10.3390/plants14193012>
- Zhang L., Kang J., Xie Q., Gong J., Shen H., Chen Y., Chen G., Hu Z. 2020. The basic helix-loop-helix transcription factor bHLH95 affects fruit ripening and multiple metabolisms in tomato. Journal of Experimental Botany 71(20): 6311-6327. <https://doi.org/10.1093/jxb/eraa363>

HOW TO CITE THIS ARTICLE

Shiri Y., Mohkami Z., Khammari I. 202x. Gene Expression Analysis of *IND*, *ALC*, *RPL*, and *FUL* in Reducing Seed Shattering in Flixweed (*Descurainia sophia*). *Agrotechniques in Industrial Crops* x(x): xx-xx. [10.22126/ATIC.2026.12895.1240](https://doi.org/10.22126/ATIC.2026.12895.1240)